9-1-2016

The Effect of Occupational Stress on Participant Success in a Worksite Weight Loss Program

Brittany Miller

Bowling Green State University, bwynn@bgsu.edu

Courtney Klebe

Bowling Green State University, cklebe@bgsu.edu

Mary-Jon Ludy

Bowling Green State University, mludy@bgsu.edu

Robin M. Tucker

Michigan State University, tucker98@msu.edu

Follow this and additional works at: https://scholarworks.bgsu.edu/publ_allied_health_pub

How does access to this work benefit you? Let us know!

Repository Citation

Miller, Brittany; Klebe, Courtney; Ludy, Mary-Jon; and Tucker, Robin M., "The Effect of Occupational Stress on Participant Success in a Worksite Weight Loss Program" (2016). *Public and Allied Health Faculty Publications*. 15.

https://scholarworks.bgsu.edu/publ_allied_health_pub/15

This Presentation is brought to you for free and open access by the Public and Allied Health at ScholarWorks@BGSU. It has been accepted for inclusion in Public and Allied Health Faculty Publications by an authorized administrator of ScholarWorks@BGSU.

The Effect of Occupational Stress on Participant Success in a Worksite Weight Loss Program

Authors: Miller B, Klebe C, Ludy MJ, Tucker RM

- Brittany Miller bwynn@bgsu.edu
- Courtney Klebe cklebe@bgsu.edu
- Mary-Jon Ludy mludy@bgsu.edu
- Robin M Tucker tucker98@msu.edu

Learning Objective: The participant will be able to identify the role of supervisor support on employee success in a weight loss program.

Background: Occupational stress has been associated with body mass index and waist circumference, but the role of occupational stress on participant outcomes at a worksite wellness weight loss program has not been examined.

Methods: 16 university faculty and staff members (18.8% faculty, 50% staff, 12.5% administration, 18.8% other) participated in a 16-week weight loss program that consisted of 7 educational sessions on nutrition and physical activity. Participants had weight, height, body mass index (BMI), body fat percentage (BF%), blood pressure (BP), total cholesterol (TC), high density lipoprotein (HDL), triglycerides (TG), and blood glucose (BG) measured before and after the program. A validated job stress survey was completed by participants at the beginning of the program, at week 10, and at the conclusion of the program.

Results: Participants lost $4.3 \text{ kg} \pm 4.8$. Significant improvements were noted in TG (33.3 ± 55.8) and BF% (2.4 ± 2.5). Supervisor support was found to be a significant predictor of weight loss, explaining 32.3% of the weight change variance. Co-worker and family support did not play a factor in weight loss during the program.

Conclusion: Worksite weight loss programs may see improved effectiveness by incorporating strategies designed to reduce occupational stress. Special attention should be placed on improving the employee-supervisor relationship to enhance likelihood of weight loss success.

Funding Source: Bowling Green State University

The Effect of Occupational Stress on Participant Success in a Worksite Weight Loss Program

Brittany Miller¹, BS; Courtney Klebe¹, MFN, RDN; Mary-Jon Ludy¹, PhD, RDN, FAND; Robin M. Tucker², PhD, RD

¹Department of Public and Allied Health, Bowling Green State University, Bowling Green, OH

²Department of Food Science and Human Nutrition, Michigan State University, East Lansing, MI

ABSTRACT

Background: Occupational stress has been associated with body mass index and waist circumference, but the role of occupational stress on participant outcomes at a worksite wellness weight loss program has not been examined.

Methods: 16 University faculty and staff members (18.8% faculty, 50.0% staff, 12.5% administration, 18.8% other) participated in a 16-week weight loss program that consisted of 7 educational sessions on nutrition and physical activity. Participants had weight, height, body mass index (BMI), body fat percentage (BF%), blood pressure (BP), total cholesterol (TC), high density lipoprotein (HDL), triglycerides (TG), and blood glucose (BG) measured before and after the program. A validated job stress survey was completed by participants at the beginning of the program, at week 10, and at the conclusion of the program.

Results: Participants lost 4.3 kg $\pm$ 4.8. Significant improvements were noted in TG (33.3 $\pm$ 55.8) and BF% (2.4 $\pm$ 2.5). Supervisor support was found to be a significant predictor of weight loss, explaining 32.3% of the weight change variance. Co-worker and family support did not play a factor in weight lost during the program.

Conclusions: Worksite weight loss programs may see improved effectiveness by incorporating strategies designed to reduce occupational stress. Special attention should be placed on improving the employee-supervisor relationship to enhance likelihood of weight loss success.

INTRODUCTION

Occupational stress can contribute to obesity and increased body mass index (BMI); consumption of energy dense, high fat, high carbohydrate foods; and decreased physical activity.^{1,2}

The role of occupational stress on participant outcomes at a worksite wellness weight loss program has not been examined.

Research Question: Does occupational stress affect weight loss in individuals attempting to lose weight in a worksite wellness weight loss program?

Hypothesis: Higher levels of occupational stress will have a negative impact on weight loss.

METHODS

Participant Characteristics

Sex	81.3% female
Race	86.7% white
Occupation Type	50.0% staff

Participant Occupations

Overall Design

*Validated NIOSH Generic Job Stress Questionnaire

METHODS (Cont.)

Educational Sessions

1: Portion Control & Individualized Energy Recommendations
2: Low-Carb vs. Low-Fat, Which is Best?
3: Eating on the Run: Fast Food Pitfalls
4: Role of Exercise in Weight Loss Plan
5: But the Internet said.... Myths about Weight Loss
6: Stress Management and Mindful Eating
7: Preventing Weight Loss Plateaus

RESULTS & DISCUSSION

Biochemical & Anthropometric Measures: Pre- & Post-Program

	Pre-Program	Post-Program
Total Cholesterol (mg/dL)	181.1 $\pm$ 32.1	190.7 $\pm$ 32.7
High-Density Lipoprotein (mg/dL)	47.8 $\pm$ 11.8	54.5 $\pm$ 15.9
Triglycerides (mg/dL)	142.25 $\pm$ 90.3	109.0 $\pm$ 70.1
Blood Glucose (mg/dL)	77.69 $\pm$ 21.3	90.0 $\pm$ 10.4
Systolic Blood Pressure (mmHg)	116.8 $\pm$ 11.8	111.7 $\pm$ 10.3
Diastolic Blood Pressure (mmHg)	76.9 $\pm$ 10.6	73.0 $\pm$ 5.4
Weight (kg)	84.7 $\pm$ 21.5	80.4 $\pm$ 17.5
Body Fat (%)	41.9 $\pm$ 6.8	39.5 $\pm$ 6.0

Biochemical & Anthropometric Measures: Mean Change

	Mean Change	SD	P-Value
Total Cholesterol (mg/dL)	↑ 9.6	22.0	0.102
High-Density Lipoprotein (mg/dL)	↑ 6.8	7.5	0.003*
Triglycerides (mg/dL)	↓ 33.3	55.8	0.031*
Blood Glucose (mg/dL)	↑ 12.3	20.8	0.032*
Systolic Blood Pressure (mmHg)	↓ 5.1	11.1	0.094
Diastolic Blood Pressure (mmHg)	↓ 3.9	7.6	0.065
Weight (kg)	↓ 4.3	4.7	0.003*
Body Fat (%)	↓ 2.4	2.5	0.002*

RESULTS & DISCUSSION (Cont.)

Predictors of Weight Change

Three forms of social support were evaluated to determine effects on weight loss: family, co-worker, and supervisor support.

- Baseline weight was the best predictor of weight change.
- Supervisor support was the best predictor of baseline weight.
- Supervisor support was significantly better at predicting weight change compared to the mean value of weight change.
- While 32.3% weight change variance was explained by supervisor support, the addition of supervisor support did not significantly improve the ability to predict weight change.
- For every one point improvement in the supervisor support score, participants lost 1.4 kg.

CONCLUSIONS

- This pilot study suggests that the employee-supervisor relationship should be considered in workplace weight loss programs. A larger sample size will help to resolve remaining questions.
- While one session of the program addressed stress management, this advice centered on how to manage overeating while stressed.
- Future participants may benefit from programming that discusses how to deal with difficult people, especially supervisors, in the work place.

REFERENCES

- Kivimaki M, Head J, Ferrie JE, Shipley MJ, Brunner E, Vahtera J, Marmot MG. Work stress, weight gain and weight loss: evidence for bidirectional effects on job strain on body mass index in the Whitehall II study. *International Journal of Obesity*. 2016;30:982-987.
- Christaki E, Kokkinos A, Costarelli V, Alexopoulos EC, Chrousos GP, Darviri C. Stress management can facilitate weight loss in Greek overweight and obese women: a pilot study. *J Hum Nutr Diet*. 2013;26(1):132-139.