

Journal of Sports Medicine and Allied Health Sciences: Official Journal of the Ohio Athletic Trainers Association

Volume 1
Issue 1 *Ohio Athletic Trainers' Association
Annual Symposium Accepted Abstracts*

Article 6

May 2015

The Anterolateral Ligament in 41 Paired Human Cadavers

Trevor Magnotti
tm013612@ohio.edu

Follow this and additional works at: <https://scholarworks.bgsu.edu/jsmahs>



Part of the [Body Regions Commons](#), [Musculoskeletal System Commons](#), and the [Sports Sciences Commons](#)

[How does access to this work benefit you? Let us know!](#)

Recommended Citation

Magnotti, Trevor (2015) "The Anterolateral Ligament in 41 Paired Human Cadavers," *Journal of Sports Medicine and Allied Health Sciences: Official Journal of the Ohio Athletic Trainers Association*: Vol. 1: Iss. 1, Article 6.

DOI: <https://doi.org/10.25035/jsmahs.01.01.06>

Available at: <https://scholarworks.bgsu.edu/jsmahs/vol1/iss1/6>

This Professional/Faculty Abstract is brought to you for free and open access by the Journals at ScholarWorks@BGSU. It has been accepted for inclusion in *Journal of Sports Medicine and Allied Health Sciences: Official Journal of the Ohio Athletic Trainers Association* by an authorized editor of ScholarWorks@BGSU.

The Anterolateral Ligament in a 41 Paired Human Cadaver

Trevor D. Magnotti, MS, AT Seth L. Faulkner, MA, AT, and Bentley A. Krause, PhD, AT

Ohio University, School of Applied Health Sciences and Wellness

Context: The anterolateral ligament of the knee has recently received significant attention in anatomical studies. The anterolateral ligament has been identified in up to 97% of unpaired cadaveric knees. This ligament is believed to assist in limiting excessive internal rotation of the knee, which could make it an important accessory stabilizer. Segund fractures, an avulsion of the distal anterolateral ligament insertion, has been implicated in 75% of anterior cruciate ligament ruptures. However, the anterolateral ligament has not been identified in bilateral knees.

Objective: To identify the anterolateral ligament in the paired knees of cadavers.

Design: A descriptive study design was used.

Setting: Medical School Cadaver Laboratory.

Patients or Other Participants: Bilateral knees of 41 bodies (n=82 knees); female = 13, mean age = 80.9 years; male = 28, mean age = 77.3 years) were dissected to confirm the presence of the anterolateral ligament and its proximal and distal attachments.

Interventions: The attachments of the anterolateral ligament were identified by gross dissection in 41 knees bilaterally. The approach of each dissection was to visualize the anterolateral ligament attachment sites from the anteromedial lip of the lateral femoral epicondyle proximally to the anterolateral aspect of the tibial condyle distally. The anterolateral ligament demonstrated an oblique, anteroinferior

course from the femoral site just proximal to the popliteal tendon to the anterolateral aspect of the proximal tibia, superior to a horizontal line connecting the fibular head and Gerdy's tubercle. The anterolateral ligament was shown to insert with the coronary ligament and the tibiofemoral joint capsule to the anterior horn of the lateral meniscus and enveloped the inferior lateral geniculate artery. Its attachment on the anterolateral tibia was found to be distinct from that of the iliotibial band.

Main Outcome Measures: The presence or absence of the anterolateral ligament was determined in paired cadavers. Once identified, the knee was internally rotated at 0° and 90° of knee flexion to confirm the integrity of the ALL.

Results: The anterolateral ligament was identified in 81 of 82 (97.561%) of knees and was present bilaterally in each of the 13 female specimens (26 knees) and in all but one knee of the male cadavers (55/56) or 98.2%.

Conclusions: Our data confirm that the anterolateral ligament is a distinct capsuloligamentous band prevalent in the majority of human knees. The anterolateral ligament insertion may be linked to Segund fractures, commonly associated with anterior cruciate ligament rupture. This anterolateral ligament should not be overlooked in the physical examination and diagnosis of rotary instabilities of the knee and is worthy of further biomechanical study.

Key Words: *Knee biomechanics; posterolateral corner; anterior cruciate ligament*