9-1-2016

The Influence of Modest Weight Gain on Taste and Smell Acuity in College Freshmen

P. Tomko
Bowling Green State University

J. Mansperger
Bowling Green State University

R. M. Tucker
Bowling Green State University

Mary-Jon Ludy
Bowling Green State University, mludy@bgsu.edu

Follow this and additional works at: https://scholarworks.bgsu.edu/publ_allied_health_pub

How does access to this work benefit you? Let us know!

Repository Citation

Tomko, P.; Mansperger, J.; Tucker, R. M.; and Ludy, Mary-Jon, "The Influence of Modest Weight Gain on Taste and Smell Acuity in College Freshmen" (2016). *Public and Allied Health Faculty Publications*. 16.
https://scholarworks.bgsu.edu/publ_allied_health_pub/16

This Presentation is brought to you for free and open access by the Public and Allied Health at ScholarWorks@BGSU. It has been accepted for inclusion in Public and Allied Health Faculty Publications by an authorized administrator of ScholarWorks@BGSU.

The Influence of Modest Weight Gain on Taste and Smell Acuity in College Freshmen

Authors: Tucker RM, Ludy MJ, Tomko PM, Mansperger J

Background: Reduced taste and smell acuities are common among those with high body mass index (BMI). Evidence of improved taste sensitivity after modest weight loss (~3-4 kg) exists in overweight individuals. Whether taste and smell decrements precede or result from weight gain is unknown. This study's purpose was to determine the effect of modest weight gain on taste and smell acuity.

Methods: College freshmen (n=60), a cohort at risk for modest weight gain, were assessed upon arrival to campus and 5 months later. Weight, BMI, and body fat percent (%BF) were measured. Intensity ratings for graded, suprathreshold concentrations of sweet (n=3) and salty (n=3) solutions, as well as a suprathreshold orange odor were assessed. Paired samples t-tests were used to indicate change from baseline to follow-up. Pearson correlation coefficients were used to identify associations between variables.

Results: Sixty participants (82% female, 92% white, 18.1 ± 0.3 years) completed testing. BMI (22.9 ± 3.8 to $23.6 \pm 4.1 \text{ kg/m}^2$) and %BF (24.5 ± 9.6 to $27.1 \pm 9.1\%$) increased at follow-up ($p < 0.001$ for both). Weight gain (62.8 ± 12.8 to $64.6 \pm 13.7 \text{ kg}$, range -2.9 to +9.1kg) occurred in 79% of participants. No changes in taste intensity for low, medium, and high concentrations of sweet or salty solutions were observed. No change in smell intensity was observed over time, although smell intensity at follow-up was negatively associated with BMI ($r = -0.36$, $p = 0.048$).

Conclusions: Modest weight gain in college freshmen did not influence taste acuity. Lack of change in intensity ratings may relate to use of suprathreshold concentrations rather than detection thresholds, or recruitment of a predominantly lean population.

Learning Outcome: Participants will be able to discuss the influence of weight change on taste and smell acuity.

The Influence of Modest Weight Gain on Taste and Smell Acuity in College Freshmen

Patrick M. Tomko,¹ Josie A. Mansperger,¹ Robin M. Tucker², Mary-Jon Ludy¹

¹Department of Public and Allied Health, Bowling Green State University; ²Department of Food Science and Human Nutrition, Michigan State University

Abstract

Background: Reduced taste and smell acuities are common among those with high BMI. Evidence of improved taste sensitivity after modest weight loss (~3-4 kg) exists in overweight individuals. Whether taste and smell decrements precede or result from weight gain is unknown. This study's purpose was to determine the effect of modest weight gain on taste and smell acuity.

Methods: College freshmen (n=60), a cohort at risk for modest weight gain, were assessed upon arrival to campus and 5 months later. Weight, BMI, and body fat percent (%BF) were measured. Intensity ratings for graded, suprathreshold concentrations of sweet (n=3) and salty (n=3) solutions, as well as a suprathreshold orange odor were assessed. Mann-Whitney-U tests were used to indicate change from baseline to follow-up. Spearman correlation coefficients were used to identify associations between variables.

Results: Sixty participants (82% female, 92% white, 18.1±0.3 years) completed testing. BMI (22.9±3.8 to 23.6±4.1 kg/m²) and %BF (24.5±9.6 to 27.1±9.1%) increased at follow-up (p<0.001 for both). Weight gain (62.8±12.8 to 64.6±13.7 kg, range -2.9 to +9.1 kg) occurred in 79% of participants. No changes in taste intensity for low, medium, and high concentrations of sweet or salty solutions were observed. No change in smell intensity was observed over time, although smell intensity at follow-up was negatively associated with BMI (r=-0.36, p=0.048).

Conclusions: Modest weight gain in college freshmen did not influence taste acuity. Lack of change in intensity ratings may relate to use of suprathreshold concentrations rather than detection thresholds, or recruitment of a predominantly lean population.

Learning Outcome: Participants will be able to discuss the influence of weight change on taste and smell acuity.

Introduction

- Taste and smell contribute to the hedonic value (or pleasantness) of food.
 - Hedonic value plays important role in food selection and intake (1).
- Individuals with high BMI typically experience impaired taste and smell sensitivity (2, 3).
 - This could cause them to eat more to achieve sufficient levels of reward (3).
- After weight loss, taste sensitivity improves (4, 5).
 - No studies have measured sensitivity changes as weight is gained.

Research Question

Does weight gain impair taste and/or smell acuity?

We hypothesized that first-year students who gained the most weight would experience decreases in taste and smell sensitivity.

Methods

Figure 1. Study protocol

- SPSS (v 24.0, IBM Corp.) was used for analysis.
- Data were not normally distributed, so non-parametric tests were used.
- Due to the large degree of variation in weight change, participants were split into quartiles.
 - Comparisons were made between the lowest (≤ 25th percentile) and highest (≥ 75th percentile) gainers.

Results

- Among all participants (n=60)
 - Weight increased 1.9±2.3 kg from August/September to January (p<0.001, range -2.9 to +9.1 kg).
 - %BF increased 1.9±3.4% from August/September to January (p<0.001, range -5.0 to +9.2%).
 - No differences in weight or %BF change were observed between males and females.
 - No consistent correlations were seen with weight gain and taste or smell intensity ratings.

Age	18.1±0.3 years
Sex	82% female (n=49)
Race	92% white (n=55)
BMI	22.9±3.8 → 23.6±4.1 kg/m ²
%BF	24.5±9.6 → 27.1±9.1%
Weight	62.8±12.8 → 64.6±13.7 kg

Table 1. Participant Characteristics

Figure 1. For the highest weight gainers (≥ 75th percentile) there were significant increases in both weight (B) and %BF (D) from baseline to follow-up (p<0.05). No significant differences were noted for the lowest weight gainers (A, C). N=15 per group and * denotes a significant difference.

Results (continued)

Figure 2. No differences in sweet intensity (ratings based on 100 mm) between individuals in the 25th percentile of weight gain compared to the 75th percentile (mean ± SD, n=15 per group). No differences were observed between baseline and follow-up for either group – indicating stability over time.

Figure 3. No differences in salt intensity (ratings based on 100 mm) between individuals in the 25th percentile of weight gain compared to the 75th percentile (mean ± SD, n=15 per group). No differences were observed between baseline and follow-up for either group – indicating stability over time.

Conclusions

- Modest weight, BMI, and %BF change did not influence perceived intensity of sweet and salty tastes or olfactory sensitivity in college freshmen.
 - This could be due to:
 - Suprathreshold concentrations rather than detection thresholds,
 - Predominantly lean rather than overweight/obese population,
 - Or the small sample size.

References & Acknowledgements

- Schutz HG, Wahl OL. Consumer perception of the relative importance of appearance, flavor and texture to food acceptance. In: Criteria of Food Acceptance Symposium Proceedings. 1981. p. 97–116.
- Richardson BE, Vander Woude EA, Sudan R, Thompson JS, Leopold DA. Altered olfactory acuity in the morbidly obese. *Obes Surg*. 2004;14: 967–969.
- Stewart JE, Keast RSJ. Recent fat intake modulates fat taste sensitivity in lean and overweight subjects. *Int J Obes*. 2012;36: 834–842.
- Umabaki M, Tsuzaki K, Kotani K, Nagai N, Sano Y, Matsuo Y, Kitaoka K, Okami Y, Sakane N, Higashi A. The improvement of sweet taste sensitivity with decrease in serum leptin levels during weight loss in obese females. *Tohoku J Exp Med*. 2010;220(4):267–71.
- Burge JC, Schaumburg JZ, Choban PS, DiSilvestro RA, Flancbaum L. Changes in patients' taste acuity after Roux-en-Y gastric bypass for clinically severe obesity. *J Am Diet Assoc*. 1995 Jun 30;95(6):666–70.

This study was supported by the DuPont Nutrition & Health/Academy of Nutrition & Dietetics Research DPG Pilot Grant Program (RMT, MJL), the USDA National Institute of Food and Agriculture (RMT), and Michigan AgBioResearch (RMT).