Journal of Sports Medicine and Allied Health Sciences: Official Journal of the Ohio Athletic Trainers Association

Volume 2
Issue 1 *Ohio Athletic Trainers' Association
Supplementary Edition*

Article 17

May 2016

Risk Factors for Concussion in Collegiate NCAA Division II Men's and Women's Soccer Athletes

Isaac S. Beckler
Cedarville University, ibeckler@cedarville.edu

Michael S. Weller
Cedarville University, mweller@cedarville.edu

Follow this and additional works at: <https://scholarworks.bgsu.edu/jsmahs>

Part of the [Sports Sciences Commons](#)

[How does access to this work benefit you? Let us know!](#)

Recommended Citation

Beckler, Isaac S. and Weller, Michael S. (2016) "Risk Factors for Concussion in Collegiate NCAA Division II Men's and Women's Soccer Athletes," *Journal of Sports Medicine and Allied Health Sciences: Official Journal of the Ohio Athletic Trainers Association*: Vol. 2: Iss. 1, Article 17.

DOI: <https://doi.org/10.25035/jsmahs.02.01.17>

Available at: <https://scholarworks.bgsu.edu/jsmahs/vol2/iss1/17>

This Undergraduate Student Abstract is brought to you for free and open access by the Journals at ScholarWorks@BGSU. It has been accepted for inclusion in *Journal of Sports Medicine and Allied Health Sciences: Official Journal of the Ohio Athletic Trainers Association* by an authorized editor of ScholarWorks@BGSU.

Cedarville University

Isaac Beckler, ATS

Michael S. Weller, M.S., AT, ATC

RISK FACTORS FOR CONCUSSION IN COLLEGIATE NCAA DIVISION II MEN'S AND WOMEN'S SOCCER ATHLETES

Objective: To identify risk factors for concussion among men's and women's college soccer athletes, and determine the likelihood of concussion based on those risk factors.

Design and Setting: A short electronic survey was sent out to NCAA Division II universities in Ohio. Survey questions included demographics, soccer-related questions, and concussion-related questions.

Participants: Men's and women's soccer athletes from division II universities in Ohio participating in varsity or junior varsity soccer at their respective universities who were at least 18 years of age.

Intervention: An online survey.

Main Outcome Measurement: Number of concussions sustained.

Results: Of the concussions sustained while playing soccer, 29.4% of female participants sustaining a concussion compared to only 17.4% male participants. Concussions occurred during a competitive match in 8 of the 9 concussions compared to during a practice. There were 2.8 concussions per 100 years for strikers as well as midfielders, while no concussions occurred while playing defender. Varsity athletes showed a high incidence of concussions (33.3%) compared to JV/Reserve athletes in which there were no concussions while playing soccer ($p = 0.018$). Among players with a concussion history, 5 of the 12 (41.7%) sustained multiple concussions. The most common MOI was contact with another player's body (36.8%). Of the 40 respondents, there were only 3 players who wore mouth guards, and 1 who wore headgear. There was no significant difference between protective equipment worn and a diagnosis of concussion ($p = .157$)

Conclusions: Being a varsity athlete, playing in a game, having a previous history of concussion, and playing as a striker or midfielder were all risk factors for concussion.

Keywords: concussion, soccer, risk factors.