Journal of Sports Medicine and Allied Health Sciences: Official Journal of the Ohio Athletic Trainers Association

Volume 3
Issue 1 *Ohio Athletic Trainers' Association*
Special Edition

Article 1

May 2017

The Relationship of Gymnastics Experience and Hand Placement Technique on Peak Ground Reaction Forces Through the Elbow

Saskia Richter
The Ohio State University, richter.sask@gmail.com

Follow this and additional works at: <https://scholarworks.bgsu.edu/jsmahs>

Part of the [Biomechanics Commons](#), [Exercise Science Commons](#), [Motor Control Commons](#), [Other Kinesiology Commons](#), [Rehabilitation and Therapy Commons](#), [Sports Medicine Commons](#), and the [Sports Sciences Commons](#)

[How does access to this work benefit you? Let us know!](#)

Recommended Citation

Richter, Saskia (2017) "The Relationship of Gymnastics Experience and Hand Placement Technique on Peak Ground Reaction Forces Through the Elbow," *Journal of Sports Medicine and Allied Health Sciences: Official Journal of the Ohio Athletic Trainers Association*: Vol. 3: Iss. 1, Article 1.

DOI: <https://doi.org/10.25035/jsmahs.03.01.01>

Available at: <https://scholarworks.bgsu.edu/jsmahs/vol3/iss1/1>

This Professional/Faculty Abstract is brought to you for free and open access by the Journals at ScholarWorks@BGSU. It has been accepted for inclusion in Journal of Sports Medicine and Allied Health Sciences: Official Journal of the Ohio Athletic Trainers Association by an authorized editor of ScholarWorks@BGSU.

The Relationship of Gymnastics Experience and Hand Placement Technique on Peak Ground Reaction Forces Through the Elbow

Saskia D. Richter, MS, ATC*, John H. Bolte, PhD*, Laura C. Boucher, PhD, ATC‡

*Division of Athletic Training, College of Medicine, ‡Division of Athletic Training, School of Health and Rehab Sciences, *‡The Ohio State University

CONTEXT

Gymnastics has a lifetime cumulative injury rate of 58%, with 13-53% of injuries occurring at the upper limb.¹⁻³ Back handsprings (BHS) are a common skill used in gymnastics.³ BHS and similar upper limb loading skills may occur over 100 times per practice.⁴ Current literature states, a turned-in hand position reduces peak ground reaction force at impact on floor, thus reducing the risk of injury.^{5,6} To our knowledge no studies analyzing the role of gymnastics experience and hand placement during a BHS on the balance beam (BB) exist.

OBJECTIVE

The purpose of this study was to identify how gymnastics experience and hand placement during a BHS on BB affects peak ground reaction forces.

DESIGN

Cross-sectional.

SETTING

Two gymnastics facilities.

PARTICIPANTS

18 competitive female gymnasts with height (cm): 153.7 $\pm$ 8.3; mass (kg): 43.1 $\pm$ 7.8; age (years): 12.7 $\pm$ 1.5. Inclusion criteria included ability to compete a BHS on BB and no practice limitations due to injury.

INTERVENTIONS

Subjects completed a demographic survey, anthropometric measurements, and 3 BHS on a BB. Hand placement technique, skill height (vertical), skill length (horizontal), angle of impact, and peak ground reaction force at impact were analyzed using high-speed cameras and portable force plates. Video analysis was completed using Dartfish Champions Software (Dartfish, Alpharetta, GA).

MAIN OUTCOME MEASURES

Gymnastics experience was quantified by years participating in gymnastics. Hand placements were grouped into categories in relation to parallel axis of the BB; neutral (both hands parallel), turned-out (one hand > 120°), turned-in (one hand < 60°), and "other". Analysis of variance (ANOVA) and pair-wise comparisons were used to identify if experience and hand placement contribute to increased force at initial hand contact.

RESULTS

Mean gymnastics experience was 7.5 $\pm$ 3.5 years. No significant difference ($p=0.58$) was found in years of experience and peak reaction force during a BHS on BB. Significance was found between hand placement and peak reaction force ($F_{(3,27)} = 15.62$; $P < .001$). Turned-in hand placement had significantly higher peak reaction force compared to all other groups ($P < .05$). BHS height (m) and length (m) was scaled from standing height (m). Mean skill height: 0.61

± 0.06 m and skill length: 0.58 ± 0.1 m. Angle of impact: $93^\circ \pm 5.8^\circ$. Mean peak vertical reaction force (N) was normalized to mass (kg): 2.5 ± 0.5 N/kg. Correlation coefficients found poor correlations between skill height, length, and angle of impact in relation to peak reaction force.

CONCLUSIONS

Results of this study suggest gymnastics experience has minimal effect on hand placement technique and peak ground reaction force upon initial impact during a BHS. Hands in a turned-in position upon landing significantly increased forces through the upper extremity, which is contrary to technique recommendations for floor exercise. Based on these data, proper hand placement technique may require additional emphasis to try to reduce forces transmitted through the upper extremity upon impact potentially reducing the risk of injury.

REFERENCES

1. O'Kane JW, Levy MR, Pietila KE, Caine DJ, Schiff MA. Survey of Injuries in Seattle Area Levels 4 to 10 Female Club Gymnasts. *Clin J Sport Med.* 2011;21(6):486-492.
2. Caine D, Harringe ML. PART 4 Chapter 10 Epidemiology of injury in gymnastics. 2013.
3. Kerr ZY, Hayden R, Barr M, Klossner DA, Dompier TP. Epidemiology of National Collegiate Athletic Association Women's Gymnastics Injuries, 2009-2010 Through 2013-2014. *J Athl Train.* 2015;50(8):870-878. doi:10.4085/1062-6050-50.7.02.
4. Sands W, McNeal J. Hand Position Back Handspring. *Technique.* 2006;1(March):8-9.
5. Farana R, Jandacka D, Uchytel J, Zahradnik D, Irwin G. The influence of hand positions on biomechanical injury risk factors at the wrist joint during the round-off skills in female gymnastics. *J Sports Sci.* 2016;414(March):1-6. doi:10.1080/02640414.2016.1158414.
6. Koh TJ, Grabiner MD, Weiker GG. Technique and ground reaction forces in the back handspring. *Am J Sports Med.* 1992;20(1):61-66.
<http://www.ncbi.nlm.nih.gov/pubmed/1554075>.

KEY WORDS: *back handspring, force, gymnastics, upper extremity*