Journal of Sports Medicine and Allied Health Sciences: Official Journal of the Ohio Athletic Trainers Association

Volume 3
Issue 1 *Ohio Athletic Trainers' Association*
Special Edition

Article 5

May 2017

Exposure to Catastrophic Injury Events and Burnout Among Athletic Trainers

Patricia Estock
Ohio University, pe506114@ohio.edu

Janet Simon
Ohio University, simonj1@ohio.edu

Follow this and additional works at: <https://scholarworks.bgsu.edu/jsmahs>

Part of the [Rehabilitation and Therapy Commons](#), [Sports Medicine Commons](#), and the [Sports Sciences Commons](#)

How does access to this work benefit you? Let us know!

Recommended Citation

Estock, Patricia and Simon, Janet (2017) "Exposure to Catastrophic Injury Events and Burnout Among Athletic Trainers," *Journal of Sports Medicine and Allied Health Sciences: Official Journal of the Ohio Athletic Trainers Association*: Vol. 3: Iss. 1, Article 5.

DOI: <https://doi.org/10.25035/jsmahs.03.01.05>

Available at: <https://scholarworks.bgsu.edu/jsmahs/vol3/iss1/5>

This Professional/Faculty Abstract is brought to you for free and open access by the Journals at ScholarWorks@BGSU. It has been accepted for inclusion in *Journal of Sports Medicine and Allied Health Sciences: Official Journal of the Ohio Athletic Trainers Association* by an authorized editor of ScholarWorks@BGSU.

Exposure to Catastrophic Injury Events and Burnout Among Athletic Trainers

Patricia Estock, AT, Janet E. Simon PhD, AT

Division of Athletic Training, Ohio University

OBJECTIVE

To assess the levels of burnout among athletic trainers (ATs) who have been exposed to catastrophic injury events and the coping strategies they utilize.

DESIGN and SETTING

Cross-sectional survey; online

PARTICIPANTS

We contacted 9881 certified members of the National Athletic Trainers' Association and invited them to participate in this study. One thousand seven surveys were completed for a response rate of 10.2% (male n=433; female n=572, prefer not to answer n=1). The majority of individuals worked in the college (n=399) or secondary school (n=436) settings and had 0-5 years of experience (n=382).

INTERVENTION

Participants completed an online survey to assess demographic information (age, sex, years as an AT, and work setting), provided care for a catastrophic injury event, Maslach Burnout Inventory-Human Services Survey (MBI-HSS), and Coping Inventory for Stressful Situations (CISS). Catastrophic injury event was defined by the National Center for Catastrophic Sport Injury Research as "fatalities, permanent disability injuries, serious injuries (fractured neck or serious head injury) even though the athlete has a full recovery, temporary or transient paralysis (athlete has no movement for a short time, but has a complete recovery), heat stroke due to exercise, or sudden cardiac arrest or sudden cardiac or severe cardiac disruption."

MAIN OUTCOME MEASURES

Scores on the MBI-HSS (personal accomplishment (PA), depersonalization (DP), and emotional exhaustion (EE) sub-scales) and the CISS (avoidance, task, and emotion-oriented sub-scales). The first multivariate ANOVA included the dependent variables (PA, DP, and EE scores) and the independent variable group (provided care for catastrophic injury vs. no exposure). The second multivariate ANOVA included the dependent variables (avoidance, task, and emotion-oriented scores) and the same independent variable group. Follow up univariate ANOVAs were conducted for each dependent variable if the overall multivariate ANOVA was significant. Alpha levels were set at $p < 0.05$ for all analyses.

RESULTS:

Five hundred eighteen (51.4%) ATs reported providing care for a catastrophic event and 489 (48.6%) did not. The group that reported providing care for a catastrophic injury had significantly worse PA (higher score worse) 46.05 ± 6.65 versus the group that has never provided care for a catastrophic injury 44.62 ± 7.71 ($p < 0.05$). The CISS score for the task-oriented sub-scale for ATs who had provided care for a catastrophic event was 30.44 ± 2.92 and 29.98 ± 3.49 for those who have not ($p < 0.05$); indicating a high use of task-oriented coping strategy. The emotion oriented score for those who had provided care for a catastrophic event was 14.26 ± 4.75 and 14.90 ± 4.17 for those who had not ($p < 0.05$); indicating a lower use of emotion-oriented coping strategy.

CONCLUSION

ATs who have provided care for a catastrophic injury event could be more likely to suffer from a lack of personal accomplishment and

exhibit task and emotion-oriented coping behaviors.

KEY WORDS: *copmg, Maslach Burnout, catastrophic injury*