11-2013

Probing Vibrationally Mediated Ultrafast Excited-state Reaction Dynamics With Multireference (caspt2) Trajectories

Patrick Z. El-Khoury

Saju Joseph

Igor Schapiro

Samer Gozem

Massimo Olivucci

Bowling Green State University, molivuc@bgsu.edu

See next page for additional authors

Follow this and additional works at: https://scholarworks.bgsu.edu/chem_pub

Part of the [Chemistry Commons](#)

How does access to this work benefit you? Let us know!

Repository Citation

El-Khoury, Patrick Z.; Joseph, Saju; Schapiro, Igor; Gozem, Samer; Olivucci, Massimo; and Tarnovsky, Alexander N., "Probing Vibrationally Mediated Ultrafast Excited-state Reaction Dynamics With Multireference (caspt2) Trajectories" (2013). *Chemistry Faculty Publications*. 102.
https://scholarworks.bgsu.edu/chem_pub/102

This Article is brought to you for free and open access by the Chemistry at ScholarWorks@BGSU. It has been accepted for inclusion in Chemistry Faculty Publications by an authorized administrator of ScholarWorks@BGSU.

Author(s)

Patrick Z. El-Khoury, Saju Joseph, Igor Schapiro, Samer Gozem, Massimo Olivucci, and Alexander N. Tarnovsky

Probing Vibrationally Mediated Ultrafast Excited-State Reaction Dynamics with Multireference (CASPT2) Trajectories

Patrick Z. El-Khoury,^{†,§} Saju Joseph,[†] Igor Schapiro,^{†,||} Samer Gozem,[†] Massimo Olivucci,^{*,†,‡} and Alexander N. Tarnovsky^{*,†}

[†]Department of Chemistry and the Center for Photochemical Sciences, Bowling Green State University, Bowling Green, Ohio 43403, United States

[‡]Dipartimento di Chimica, Università di Siena, via De Gasperi 2, I-53100 Siena, Italy

S Supporting Information

ABSTRACT: Excited-state trajectories computed at the complete active space second-order perturbation theory (CASPT2) reveal how vibrational excitation controls the molecular approach to the intersection space that drives the photodissociation of a prototypical halogenated methyl radical, namely CF₂I. Translating the Franck–Condon structure along the ground-state CASPT2 vibrational modes in this system followed by propagating the displaced structures in the first excited doublet state simulates specific vibrational excitations and vibrationally mediated dynamics, respectively. Three distinct situations are encountered: the trajectories (i) converge to an energetically flat segment of the intersection space, (ii) locate a segment of the intersection space, and (iii) access a region where the intersection space degeneracy is lifted to form a ridge of avoided crossings. The computational protocol documented herein can be used as a tool to design control strategies based on selective excitation of vibrational modes, including adaptive feedback schemes using coherent light sources.

INTRODUCTION

Nuclear motion is a prerequisite for chemical transformations, the control of which has rattled several generations of scientists. The key to controlling the outcome of a chemical reaction is the ability to selectively localize energy in a favorable reaction coordinate. A viable approach is to employ vibrational excitation, which may steer a molecular system toward a desired reaction pathway,¹ either in the ground or in the excited electronic states, as demonstrated in early studies.^{2–4} To date, this concept and techniques which stemmed from such studies have been applied in various systems, including (i) vibrationally mediated photodissociation and bimolecular reactions in the gas phase,^{4–10} (ii) state-selective dissociative chemisorption on metal surfaces^{11,12} and (iii) more recently, condensed-phase photochemical reactions such as cis–trans isomerization and energy transfer in natural and artificial light-harvesting complexes, where selective excitation of vibrational modes is achieved using optimal control schemes^{13,14} utilizing femto-second laser pulse shaping.^{15–20}

Efficient control over chemical reaction dynamics presupposes a molecular level understanding of complex multidimensional energy landscapes. In photochemical transformations, the picture is further complicated by molecular evolution on at least two electronic surfaces. Recognizing that the degeneracy of electronic states is a common occurrence in polyatomics,^{21–26} one also needs to identify the molecular and electronic structures of conical intersections (CIs), which mediate ultrafast radiationless decay pathways in photoinduced processes. To account for multiple electronic configurations encountered at the degeneracy points, designing predictive

schemes for vibrationally mediated photochemical processes defaults a practitioner to the use of tools of multiconfigurational quantum chemistry, i.e., wave functions flexible enough to account for multiple electronic configurations. A general description of vibrationally mediated nonadiabatic photochemical transformations also ought to account for all decay paths dynamically and energetically feasible to an evolving molecular system in multidimensional conformational subspace. The major challenge in achieving optimal control is that it requires an extensive knowledge of all such decay paths.

In this work, we model the *excited-state* branch of a vibrational-state-specific photochemical/photophysical transformation in a prototypical halogenated alkane radical, CF₂I, by computing on-the-fly excited-state adiabatic trajectories using the complete active space second-order perturbation (CASPT2)²⁷ level of theory. This is a development of our previous study of CF₂I, where we mapped the minimum energy photochemical reaction pathways from the lowest-lying electronically excited state using the CASPT2//CASSCF protocol.²⁸ A CI was located between the ground D₀ and first excited D₁ doublet states at the CASSCF level and found to drive the excited-state species to the reactant (CF₂I) and photoproduct (CF₂ + I) wells on the ground state. Trajectories based on gradients from multireference approaches have been recently computed by several groups.^{25,29–31} However, in the present study we systematically sample the effect of the degree

Received: August 22, 2013

Revised: August 31, 2013

Published: September 3, 2013

Figure 2. Left column: 2D energy plots of the D_0 and D_1 states of CF_2I as a function of the displacement along the $\Delta\nu_{11}$ and $\Delta\nu_{21}$, $\Delta\nu_{11}$ and $\Delta\nu_{31}$, and $\Delta\nu_{11}$ and $\Delta\nu_{41m}$ vectors. Right column: the corresponding $D_0 - D_1$ energy gaps. The full line on the coordinate base represents loci of the intersection space. Dashed lines indicate the region where the degeneracy has been lifted.

which is within the desired accuracy. Finally, the trajectories were initiated with zero kinetic energy. Indeed, by displacing the molecule along a vibrational mode ν_n we populate ν_n only. At the turning point where excitation occurs, the molecule has zero kinetic energy along that mode. Moreover, we assume that all other modes have zero energy to isolate the effect of only ν_n and to not bias the trajectories by our choice of nuclear velocities, but rather to allow the excited-state gradient to drive the starting structures. If one randomly populates other modes by geometric distortion or by adding velocities along these modes, it would no longer be possible to isolate the effect of the ν_n alone on the trajectory calculations. We stress that the objective of our work is to explore vibrational eigenstate-specific effects, which is why we do not adopt the standard practice of random sampling³⁶ to simulate wavepacket preparation/propagation. Our objective cannot be achieved by randomly populating all the vibrational levels at the same time, where the effect of each vibrational mode will be lost.

RESULTS AND DISCUSSION

A reference excited-state CASPT2 trajectory ν_{00} is computed by releasing the system D_0 equilibrium structure (the FC point) along the D_1 energy surface with no initial kinetic energy, and

terminated at the first intercepted CI point, which is then taken as the reference CI, $\nu_{00}\text{-CI}$. The fact that the reference trajectory hits a CI point in only 15 fs (Figure 1) indicates that the reaction coordinate must have a large component along one of the branching plane vectors²¹ at $\nu_{00}\text{-CI}$. We begin by inspecting the D_1 trajectories resulting from translating the FC point along the lowest energy normal mode, ν_1 , Figure 1a. The resulting trajectories bear a resemblance to the reference ν_{00} trajectory in terms of (i) the resulting time-dependent energy profiles, (ii) energies of the D_1 FC points, and (iii) the relative energy of the different $D_1\text{-}D_0$ CI points reached by these trajectories. Because the molecules following ν_{00} and ν_1 trajectories have different initial geometries and the CI point structures have slightly different geometries along the FCI angle, the above similarities can be rationalized only by the fact that ν_1 is a low-energy mode mostly displacing the trajectories along a flat region of the intersection subspace. In other words, it appears that for displacement along ν_1 , excited-state species are channeled through nearly parallel paths along the $N - 2$ -dimensional intersection subspace accessed by ν_{00} , where $N = 6$ is the number of vibrational degrees of freedom in the system.

On the other hand, the trajectories resulting from the displacement along ν_2 do not have the properties (i–iii) of

those displaced along ν_1 . When the FC point is displaced along ν_2 , a locus of conical intersections, i.e., a segment of the intersection space with different geometries and energies, is traced by the CASPT2 excited-state trajectories. This implies that ν_2 like ν_1 , has a projection on the intersection space, but samples a steeper excited state in the FC region. To support the conclusions drawn above, we scanned the D_0 and D_1 potential energy surfaces in the vicinity of the ν_{00} -CI structure. We displaced the ν_{00} -CI structure along $\Delta\nu_{11}$ and $\Delta\nu_{21}$ vectors, where $\Delta\nu_{ki}$ is defined as the difference in internal coordinates between ν_{00} and ν_{ki} , and where the subscript k is the vibrational mode number and the unity designates the displacement of 0.1 au, Figure 2a,b. We found that the displacement of the ν_{00} -CI along $\Delta\nu_{11}$ causes little degeneracy splitting, whereas the splitting is even less pronounced along $\Delta\nu_{21}$, and a combination of $\Delta\nu_{11}$ and $\Delta\nu_{21}$ leads to a plane of intersection points. Therefore, populating a normal mode that has a large projection on the space orthogonal to the branching plane defined by the gradient difference and derivative coupling vectors allows sampling of different conical intersection points along the seam, Figure 1a,b.

We then translate the FC point along ν_3 , a displacement that resembles the excited-state reaction vector pointing toward the minimum energy conical intersection characterized in the CASSCF study.²⁸ Not surprisingly, the Franck–Condon region is steep for translation along ν_3 , as is evident from inspecting Figure 1c. As a result, a small displacement along this mode causes (i) a large drop in the energy of the starting structure, and (ii) a systematic decrease in the relative energies of the intercepted CIs for larger displacements. Nonetheless, the computed trajectories reach their respective degeneracy points at the same time. This is interpreted in terms of cancellation effects. Namely, a decrease in acceleration due to smaller local curvature of PES is annulled by a smaller “time of travel” from the ν_3 -displaced starting structure to the seam. Therefore, ν_3 is interpreted as orthogonal to the intersection space and parallel to one of the branching plane vectors such as to the gradient different vector, i.e., the difference in the gradients of the D_1 and D_0 energy surfaces.

A conceptually different and practically interesting scenario is encountered when the system is displaced along ν_4 , Figure 1d. The positive ν_4 displacement propels CF_2I toward the CI seam. In contrast, the negative ν_4 displacement leads to a region where D_0 – D_1 degeneracy is gradually lifted. In fact, these computed trajectories trace an avoided crossing and the prepared excited-state species do not reach the seam within the computed time scale. The additional time spent by the excited-state species at the avoided crossing would be expected to slave the ν_4 populated species to intra- and intermolecular vibrational energy dissipation pathways in practical applications. On the basis of the large energy drop of the starting structure along ν_4 , this mode must have a projection on the reaction coordinate vector, similar to ν_3 . We note that the reaction vector has a large component of C–I stretching.²⁸ In fact, ν_3 , ν_4 , and to a smaller extent ν_2 , have a significant C–I stretch character (Figure 1b–d), which explains their dynamic similarities. Consistent with our prior analysis in Figure 2a,b, the topological features resulting from the ν_{00} -CI structure displaced along $\Delta\nu_{31}$ and $\Delta\nu_{41}$ vectors in combination with $\Delta\nu_{11}$ reveal that both $\Delta\nu_{31}$ and $\Delta\nu_{41}$ exhibit a surface with a conical shape and hence have components along the branching plane, supporting our reasoning above, Figure 2c–f. Con-

versely, ν_1 and ν_2 appear to have a larger projection on the intersection space.

CONCLUSIONS

Overall, uncovering the relationship between the normal modes, the topologies of the ground and excited states, and the branching plane vectors, aided either (i) by vibrational mediation where the laser frequency is simply tuned to selectively populate specific vibrational modes or (ii) by creating superpositions of vibrational states in optimal control schemes and nonlinear spectroscopy, is prerequisite for understanding how effective control over the excited-state dynamics and, therefore, the outcome of a nonadiabatic photochemical reaction can be achieved. Clearly, controlling the excited-state path is a challenge but the only sensible approach is to learn about it as much as possible and computational theory helps us to do that. We rigorously illustrate how excited-state species access different regions of the conical intersection seam or avoid it by using on-the-fly CASPT2 excited-state dynamics of a small polyatomic radical, which trace the locus of the CI in the basis of the vibrational normal modes, thus enabling us to visualize different segments of the intersection space. The concepts illustrated herein, combined with ultrafast time-resolved techniques and the advances in multidimensional quantum simulations with an explicit inclusion of the laser field,³⁷ should aid in designing and rationalizing the outcome of experiments aiming at exploiting vibrational-state specificity to optimize photochemical reactivity.

ASSOCIATED CONTENT

Supporting Information

Computational details, energy conservation plot, active space orbitals, energy profiles, Cartesian coordinates, and movies illustrating the molecular motion for four trajectories in which the FC structure is displaced along ν_1 – ν_4 . This material is available free of charge via the Internet at <http://pubs.acs.org>.

AUTHOR INFORMATION

Corresponding Authors

*A. N. Tarnovsky: e-mail, atarnov@bgsu.edu.

*M. Olivucci: e-mail, molivuc@bgsu.edu, olivucci@unisi.it.

Present Addresses

[§]Physical Sciences Division, Pacific Northwest National Laboratory, P.O. Box 999, Richland, Washington 99352, United States.

^{||}MPI für Chemische Energiekonversion, Stiftstrasse 34-36, D-45470 Mülheim an der Ruhr, Germany.

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

This work was supported by the NSF (A.N.T., CAREER award, CHE-0847707; M.O., CHE-1152070). A.N.T. and P.E.K. gratefully acknowledge an allocation of computer time from the Ohio Supercomputer Center and the Extreme Science and Engineering Discovery Environment (XSEDE, TG130003).

REFERENCES

- (1) Polanyi, J. C. Some Concepts in Reaction Dynamics. *Acc. Chem. Res.* 1972, 5, 161–168.

- (2) Odiorne, T. J.; Brooks, P. R.; Kasper, J. V. V. Molecular Beam Reaction of K with HCl: Effect of Vibrational Excitation of HCl. *J. Chem. Phys.* **1971**, *55*, 1980–1982.
- (3) Schatz, G. C.; Colton, M. C.; Grant, J. L. A Quasiclassical Trajectory Study of the State-to-State Dynamics of $\text{H} + \text{H}_2\text{O} \rightarrow \text{OH} + \text{H}_2$. *J. Phys. Chem.* **1984**, *88*, 2971–2977.
- (4) Crim, F. F. Bond-Selected Chemistry: Vibrational State Control of Photodissociation and Bimolecular Reaction. *J. Phys. Chem.* **1996**, *100*, 12725–12734.
- (5) Hause, M. L.; Yoon, Y. H.; Case, A. S.; Crim, F. F. Dynamics at Conical Intersections: The Influence of O–H Stretching Vibrations on the Photodissociation of Phenol. *J. Chem. Phys.* **2008**, *128*, 104307.
- (6) Zare, R. N. Laser Control of Chemical Reactions. *Science* **1998**, *279*, 1875–1879.
- (7) Crim, F. F. Chemical Dynamics of Vibrationally Excited Molecules: Controlling Reactions in Gases and on Surfaces. *Proc. Natl. Acad. Sci. U. S. A.* **2008**, *105*, 12654–12661.
- (8) Crim, F. F. Molecular Reaction Dynamics across the Phases: Similarities and Differences. *Faraday Discuss.* **2012**, *157*, 9–26.
- (9) Yan, S.; Wu, Y.-T.; Zhang, B.; Yue, X.-F.; Liu, K. Do Vibrational Excitations of CHD_3 Preferentially Promote Reactivity Toward the Chlorine Atom? *Science* **2007**, *316*, 1723–1726.
- (10) Wang, F.; Lin, J.-S.; Liu, K. Steric Control of the Reaction of CH Stretch–Excited CHD_3 with Chlorine Atom. *Science* **2011**, *331*, 900–903.
- (11) Killelea, D. R.; Campbell, V. L.; Shuman, N. S.; Utz, A. L. Bond-Selective Control of a Heterogeneously Catalyzed Reaction. *Science* **2008**, *319*, 790–793.
- (12) Chen, L.; Ueta, H.; Bison, R.; Beck, R. D. Vibrationally Bond-Selected Chemisorption of Methane Isotopologues on Pt(111) Studied by Reflection Absorption Infrared Spectroscopy. *Faraday Discuss.* **2012**, *157*, 285–295.
- (13) Judson, R. S.; Rabitz, H. Teaching Lasers to Control Molecules. *Phys. Rev. Lett.* **1992**, *68*, 1500–1503.
- (14) Kosloff, R.; Rice, S. A.; Gaspard, P.; Tersigni, S.; Tannor, D. J. Wavepacket Dancing: Achieving Chemical Selectivity by Shaping Light Pulses. *Chem. Phys.* **1989**, *139*, 201–220.
- (15) Brixner, T.; Gerber, G. Quantum Control of Gas-Phase and Liquid-Phase Femtochemistry. *ChemPhysChem* **2003**, *4*, 418–438.
- (16) Herek, J. H.; Wohlleben, W.; Cogdell, R. J.; Zeidler, D.; Motzkus, M. Quantum Control of Energy Flow in Light Harvesting. *Nature* **2002**, *417*, 533–535.
- (17) Dietzek, B.; Bruggemann, B.; Pascher, T.; Yartsev, A. Mechanisms of Molecular Response in the Optimal Control of Photoisomerization. *Phys. Rev. Lett.* **2006**, *97*, 258301.
- (18) Prokhorenko, V. I.; Nagy, A. M.; Waschuk, S. A.; Brown, L. S.; Birge, R. R.; Miller, R. J. D. Coherent Control of Retinal Isomerization in Bacteriorhodopsin. *Science* **2006**, *313*, 1257–1261.
- (19) Silberberg, Y. Quantum Coherent Control for Nonlinear Spectroscopy and Microscopy. *Annu. Rev. Phys. Chem.* **2009**, *60*, 277–292.
- (20) Buckup, T.; Hauer, J.; Voll, J.; Vivie-Riedle, R.; Motzkus, M. A General Control Mechanism of Energy Flow in the Excited State of Polyenic Biochromophores. *Faraday Discuss.* **2011**, *153*, 213–225.
- (21) Yarkony, D. R. Nonadiabatic Quantum Chemistry - Past, Present, and Future. *Chem. Rev.* **2012**, *112*, 481–498.
- (22) Bernardi, F.; Olivucci, M.; Robb, M. A. Potential Energy Surface Crossings in Organic Photochemistry. *Chem. Soc. Rev.* **1996**, *25*, 321–328.
- (23) Worth, G. A.; Cederbaum, L. S. Beyond Born-Oppenheimer: Molecular Dynamics through a Conical Intersection. *Annu. Rev. Phys. Chem.* **2004**, *55*, 127–158.
- (24) Paterson, M. J.; Bearpark, M. J.; Robb, M. A.; Blancafort, L.; Worth, G. A. Conical Intersections: A Perspective on the Computation of Spectroscopic Jahn–Teller Parameters and the Degenerate ‘Intersection Space’. *Phys. Chem. Chem. Phys.* **2005**, *7*, 2100–2115.
- (25) Levine, B. G.; Martinez, T. J. Isomerization through Conical Intersections. *Annu. Rev. Phys. Chem.* **2007**, *58*, 613–634.
- (26) Matsika, S.; Krause, P. Nonadiabatic Events and Conical Intersections. *Annu. Rev. Phys. Chem.* **2011**, *62*, 621–643.
- (27) Aquilante, F.; De Vico, L.; Ferré, N.; Ghigo, G.; Malmqvist, P.-Å.; Neogrady, P.; Pedersen, T. B.; Pitonak, M.; Reiher, M.; Roos, B. O.; Serrano-Andrés, L.; Urban, M.; Veryazov, V.; Lindh, R. MOLCAS 7: The Next Generation. *J. Comput. Chem.* **2010**, *31*, 224–247.
- (28) El-Khoury, P. Z.; George, L.; Kalume, A.; Schapiro, I.; Olivucci, M.; Tarnovsky, A. N.; Reid, S. A. Matrix Isolation and Computational Studies of the CF_2I Radical. *Chem. Phys. Lett.* **2010**, *496*, 68–73.
- (29) Martinez, T. J. Insights for Light-Driven Molecular Devices from Ab Initio Multiple Spawning Excited-State Dynamics of Organic and Biological Chromophores. *Acc. Chem. Res.* **2006**, *39*, 119–126.
- (30) Barbatti, M.; Lischka, H. Nonadiabatic Deactivation of 9H-Adenine: A Comprehensive Picture Based on Mixed Quantum-Classical Dynamics. *J. Am. Chem. Soc.* **2008**, *130*, 6831–6839.
- (31) Barbatti, M.; Aquino, A. J. A.; Szymczak, J. J.; Nachtigallova, D.; Hobza, P.; Lischka, H. Relaxation Mechanisms of UV-Photoexcited DNA and RNA Nucleobases. *Proc. Natl. Acad. Sci. U. S. A.* **2010**, *107*, 21453–21458.
- (32) Pulay, P. A Perspective on the CASPT2 Method. *Int. J. Quantum Chem.* **2011**, *111*, 3273–3279.
- (33) Roca-Sanjuán, D.; Aquilante, F.; Lindh, R. Multiconfiguration Second-Order Perturbation Theory Approach to Strong Electron Correlation in Chemistry and Photochemistry. *WIREs Comput. Mol. Sci.* **2012**, *2*, 585–603.
- (34) Schapiro, I.; Ryazantsev, M. N.; Frutos, L. M.; Ferré, N.; Lindh, R.; Olivucci, M. The Ultrafast Photoisomerizations of Rhodopsin and Bathorhodopsin Are Modulated by Bond Length Alternation and HOOP Driven Electronic Effects. *J. Am. Chem. Soc.* **2011**, *133*, 3354–3364.
- (35) Griebel, M.; Knapek, S.; Zumbusch, G. *Numerical Simulation in Molecular Dynamics*; Springer: Berlin, 2007.
- (36) Pederzoli, M.; Pittner, J.; Barbatti, M.; Lischka, H. Nonadiabatic Molecular Dynamics Study of the cis-trans Photoisomerization of Azobenzene Excited to the S_1 State. *J. Phys. Chem. A* **2011**, *115*, 11136–11143.
- (37) Petersen, J.; Wohlgemuth, M.; Sellner, B.; Bonačić-Koutecký, V.; Lischka, H.; Mitrić, R. Laser Pulse Trains for Controlling Excited State Dynamics of Adenine in Water. *Phys. Chem. Chem. Phys.* **2012**, *14*, 4687–4694.