Journal of Sports Medicine and Allied Health Sciences: Official Journal of the Ohio Athletic Trainers Association

Volume 4
Issue 1 *OATA Supplemental Issue*

Article 22

May 2018

Changes in SEBT Scores in College Basketball Players Participating in a Preventative Ankle Program

Haley Potters

Jennifer Miller

Kurt Wilson

Follow this and additional works at: <https://scholarworks.bgsu.edu/jsmahs>

Part of the [Biomechanics Commons](#), [Exercise Science Commons](#), [Motor Control Commons](#), [Other Kinesiology Commons](#), [Rehabilitation and Therapy Commons](#), [Sports Medicine Commons](#), and the [Sports Sciences Commons](#)

[How does access to this work benefit you? Let us know!](#)

Recommended Citation

Potters, Haley; Miller, Jennifer; and Wilson, Kurt (2018) "Changes in SEBT Scores in College Basketball Players Participating in a Preventative Ankle Program," *Journal of Sports Medicine and Allied Health Sciences: Official Journal of the Ohio Athletic Trainers Association*: Vol. 4: Iss. 1, Article 22.

DOI: <https://doi.org/10.25035/jsmahs.04.01.22>

Available at: <https://scholarworks.bgsu.edu/jsmahs/vol4/iss1/22>

This Undergraduate Student Abstract is brought to you for free and open access by the Journals at ScholarWorks@BGSU. It has been accepted for inclusion in Journal of Sports Medicine and Allied Health Sciences: Official Journal of the Ohio Athletic Trainers Association by an authorized editor of ScholarWorks@BGSU.

Changes in SEBT Scores in College Basketball Players Participating in a Preventative Ankle Program

Haley Potters; Jennifer Miller; Kurt Wilson AT, CSCS

Division of Athletic Training, Ohio Northern University

INTRODUCTION

Balance deficits have been associated with increased ankle injury rates among athletes.¹ Preventative exercise programs have been developed to help with injury prevention, but the effectiveness of these programs has not been assessed by a test that is easily administered by clinicians.

OBJECTIVE

The purpose of this study was to use the Star Excursion Balance Test (SEBT) to determine whether a 5-week preventative exercise program would improve participants' reach distances.

SETTING

Athletic training research laboratory.

PARTICIPANTS

Six male and seven female college basketball players.

METHODS

After completing a baseline SEBT, the participants took part in a 5-week preventative exercise program which incorporated postural stability, strength and plyometric exercises. Following the program, the SEBT was repeated and the results served as the outcome measurements.

INTERVENTION

5-week preventative exercise program.

MAIN OUTCOME MEASUREMENTS

Star Excursion Balance Test

RESULTS

Following the preventative program the male and female participants both significantly increased reach distances. The female participants increased significantly in the left leg posteromedial direction (96.6 ± 3.67 to 105.67 ± 3). The male participants increased significantly in the left leg anterior (65.89 ± 2.89 to 74.38 ± 3.62), posteromedial (98.28 ± 3.81 to 108.12 ± 4.09) and posterolateral (84.69 ± 5.95 to 99.35 ± 5.57) directions.

CONCLUSION

A 5-week preventative exercise program may improve the balance of college basketball players and so reduce the likelihood of sustaining an ankle injury.

REFERENCES

1. Gribble PA, Hertel J, Plisky P. Using the Star Excursion Balance Test to Assess Dynamic Postural-Control Deficits and outcomes in Lower Extremity Injury: A literature and Systemic Review. *Journal of Athletic Training*. 2012;47(3):339-357. doi:10.4085/1062-6050-47.3.08

KEY WORDS: *balance; preventative program; SEBT; basketball players; injury prevention; postural stability; strength; plyometric.*