

Journal of Sports Medicine and Allied Health Sciences: Official Journal of the Ohio Athletic Trainers Association

Volume 6
Issue 1 *OATA 2020 Supplemental Issue*

Article 14

May 2020

Comparison of Dynamic Balance in Male and Female Collegiate Lacrosse and Soccer Athletes

Elizabeth Carter
Ohio Northern University, e-carter.2@onu.edu

Brandon Bartlome
Ohio Northern University

Kurt Wilson
Ohio Northern University

Follow this and additional works at: <https://scholarworks.bgsu.edu/jsmahs>



Part of the [Biomechanics Commons](#), [Exercise Science Commons](#), [Motor Control Commons](#), [Other Kinesiology Commons](#), [Rehabilitation and Therapy Commons](#), [Sports Medicine Commons](#), and the [Sports Sciences Commons](#)

Recommended Citation

Carter, Elizabeth; Bartlome, Brandon; and Wilson, Kurt (2020) "Comparison of Dynamic Balance in Male and Female Collegiate Lacrosse and Soccer Athletes," *Journal of Sports Medicine and Allied Health Sciences: Official Journal of the Ohio Athletic Trainers Association*: Vol. 6 : Iss. 1 , Article 14.

DOI: <https://doi.org/10.25035/jsmahs.06.01.14>

Available at: <https://scholarworks.bgsu.edu/jsmahs/vol6/iss1/14>

This Undergraduate Student Abstract is brought to you for free and open access by the Journals at ScholarWorks@BGSU. It has been accepted for inclusion in *Journal of Sports Medicine and Allied Health Sciences: Official Journal of the Ohio Athletic Trainers Association* by an authorized editor of ScholarWorks@BGSU.

Comparison of Dynamic Balance in Male and Female Collegiate Lacrosse and Soccer Athletes

Eli Carter; Brandon Bartlome; Kurt Wilson, ATC
Ohio Northern University; Human Performance and Sport Sciences, Athletic Training Department

CONTEXT

Deficits in dynamic balance have been a strong predictor of lower extremity injury. Lower extremity dynamic balance between gender and sport have been examined but there is inconclusive data that supports a difference.

OBJECTIVE

To determine a difference in dynamic balance between NCAA DIII men's and women's soccer/lacrosse using the modified star excursion balance test (SEBT).

DESIGN AND SETTING

Dynamic balance assessment using the modified SEBT. Subjects have one test and will complete 3 trials using their dominant leg. Average score of the 3 trials will be used in our analysis. Subjects will be evaluated in a controlled clinical environment on multipurpose flooring.

PARTICIPANTS

23 females and 25 males participating in varsity soccer or lacrosse at Ohio Northern University during the 2019-2020 season. Subjects are 18-23 years old and have not had a LE injury, vestibular problems or concussion in the last 12 weeks.

INTERVENTION

Subjects were shown a video instructing proper technique on SEBT prior to testing and allowed 3 practice trials prior to testing.

MAIN OUTCOME MEASUREMENT

Reach distances were measured in the anterior, posteromedial and posterolateral direction and normalized for leg length: (excursion distance/leg length x 100). Mean normalized reach distances for each gender were compared using a T-Test.

RESULTS

Male participants demonstrated a significantly greater posteromedial reach than female participants (male: 100.2 (\pm 10.6), female: 94.0 (\pm 9.03), $p=0.03$). No significant differences in posterolateral or anterior directions noted.

CONCLUSION

Among collegiate soccer and lacrosse players, male athletes appear to exhibit superior dynamic balance in the posteromedial direction when compared to females. The addition of posteromedial focused dynamic balance exercises may be beneficial to prevent lower extremity injury in female soccer and lacrosse athletes.

KEY WORDS: *SEBT, Dynamic Balance, Gender, Sport, LE Injury*