

Journal of Sports Medicine and Allied Health Sciences: Official Journal of the Ohio Athletic Trainers Association

Volume 6
Issue 1 *OATA 2020 Supplemental Issue*

Article 1

May 2020

Effect of Fatigue on Lumbopelvic-Hip Complex Muscle Activation and Lower Extremity Biomechanics

Erika Zambarano

The University of Toledo, erika.zambarano@rockets.utoledo.edu

Neal Glaviano

The University of Toledo

Lucinda Bouillon

The University of Toledo

Grant Norte

The University of Toledo

Follow this and additional works at: <https://scholarworks.bgsu.edu/jsmahs>



Part of the [Biomechanics Commons](#), [Exercise Science Commons](#), [Motor Control Commons](#), [Other Kinesiology Commons](#), [Rehabilitation and Therapy Commons](#), [Sports Medicine Commons](#), and the [Sports Sciences Commons](#)

How does access to this work benefit you? Let us know!

Recommended Citation

Zambarano, Erika; Glaviano, Neal; Bouillon, Lucinda; and Norte, Grant (2020) "Effect of Fatigue on Lumbopelvic-Hip Complex Muscle Activation and Lower Extremity Biomechanics," *Journal of Sports Medicine and Allied Health Sciences: Official Journal of the Ohio Athletic Trainers Association*: Vol. 6: Iss. 1, Article 1.

DOI: <https://doi.org/10.25035/jsmahs.06.01.01>

Available at: <https://scholarworks.bgsu.edu/jsmahs/vol6/iss1/1>

This Professional/Faculty Abstract is brought to you for free and open access by the Journals at ScholarWorks@BGSU. It has been accepted for inclusion in *Journal of Sports Medicine and Allied Health Sciences: Official Journal of the Ohio Athletic Trainers Association* by an authorized editor of ScholarWorks@BGSU.

Effect of Fatigue on Lumbopelvic-Hip Complex Muscle Activation and Lower Extremity Biomechanics

Erika Zambarano, MS, AT, ATC; Neal Glaviano, PhD, AT, ATC; Lucinda Bouillon, PT, PhD; Grant Norte, PhD, AT, ATC, CSCS

The University of Toledo; College of Health and Human Services

OBJECTIVE

Instability of the lumbopelvic-hip complex (LPHC) is linked to lower extremity injury. Fatigue has shown to alter LPHC muscle activity and function. Our purpose was to examine the effect of fatigue on LPHC stability, lower extremity biomechanics, and LPHC muscle activation.

DESIGN AND SETTING

Descriptive study.

PARTICIPANTS

Twenty-two healthy, active individuals (12 females, 10 males; 23.5±3.1yrs; 1.73±0.09m; 76.1±10.7kg).

INTERVENTION

Participants completed a Seated Trunk Control Test (STCT) involving balancing on a wobble board placed on a raised surface while the researcher counted errors occurring in 30 seconds. Participants completed the STCT and single leg squats (SLS) before and after a 35-minute full-body fatiguing protocol.

MAIN OUTCOME MEASUREMENT

LPHC stability was measured using time to first error (TTE) and number of errors (NE) in the STCT. Activity of the rectus abdominis (RA), external oblique, internal oblique (IO), erector spinae, gluteus medius vastus lateralis, and vastus medialis during the tasks was recorded using wireless electromyography (EMG) and normalized to peak activation. Frontal and sagittal plane

kinematics of the trunk, hip, and knee during SLS were assessed using 2D video analysis. Paired t-tests measured differences between pre- and post-fatigue measurements.

RESULTS

TTE significantly decreased (pre: 13.44±9.54seconds, post: 8.47±8.11seconds, p=0.03) and NE significantly increased (pre: 5.73±4.99, post: 8.30±5.47, p=0.016) when fatigued. RA activity increased in the STCT (pre: 15.31±15.73, post: 28.70±31.11, p=0.047) and SLS (pre: 7.43±6.03, post: 16.83±19.97, p=0.014) when fatigued. Knee flexion angle increased (pre: 98.7±14.2; post: 102.1±15.5, p=0.042) when fatigued, indicating less knee flexion movement during the SLS. No significant changes happened in frontal plane movement.

CONCLUSION

Full-body fatigue influences LPHC stability and activation of superficial musculature used in the task. The trunk rigidity necessary to achieve stability and complete a functional task requires increased muscle activation in a fatigued state, mimicking findings in studies evaluating walking gait after exercise. Fatigue alters muscle activity of the LPHC during a SLS; however, the impact fatigue has on the quality of movement and biomechanics should be further investigated. Clinicians can use these findings to consider training the LPHC in a fatigued state.

KEY WORDS: Core, Stability, Risk Factors