International Journal of Aquatic Research and Education

Volume 11
Number 2 *The Lifesaving Foundation
Conference 2018 Booklet of Abstracts*

Article 23

10-22-2018

The Results of the PARAMEDIC 2 Clinical Trial

John Long
RLSS, Consultant, University of Warwick (UK)

Follow this and additional works at: <https://scholarworks.bgsu.edu/ijare>

Part of the [Exercise Physiology Commons](#), [Exercise Science Commons](#), [Health and Physical Education Commons](#), [Leisure Studies Commons](#), [Other Rehabilitation and Therapy Commons](#), [Outdoor Education Commons](#), [Recreation Business Commons](#), [Sports Management Commons](#), [Sports Sciences Commons](#), [Sports Studies Commons](#), and the [Tourism and Travel Commons](#)

How does access to this work benefit you? Let us know!

Recommended Citation

Long, John (2018) "The Results of the PARAMEDIC 2 Clinical Trial," *International Journal of Aquatic Research and Education*: Vol. 11: No. 2, Article 23.

DOI: <https://doi.org/10.25035/ijare.11.02.23>

Available at: <https://scholarworks.bgsu.edu/ijare/vol11/iss2/23>

This Abstract is brought to you for free and open access by the Journals at ScholarWorks@BGSU. It has been accepted for inclusion in International Journal of Aquatic Research and Education by an authorized editor of ScholarWorks@BGSU.

The Results of the PARAMEDIC 2 Clinical Trial

John Long

Patient/Public Involvement Consultant on Trial at the Clinical Trials Unit of
the Medical School of the University of Warwick

johnpatlong@aol.com

Background

Concern about adrenaline (known as epinephrine in the USA) as a treatment for out-of-hospital cardiac arrest led to the International Liaison Committee on Resuscitation and other international organisations to call for a placebo-controlled trial to see if adrenaline is safe and effective to use as a treatment.

Trial

A randomised double blind trial involving 8014 out-of-hospital cardiac arrest patients was run over four years in five National Health Ambulance Trust areas in the United Kingdom with paramedics administering adrenaline or saline placebo along with standard CPR techniques. The trial was co-ordinated by the Clinical Trials Unit of the Medical School of the University of Warwick. Professor Gavin Perkins was the Chief Investigator. The primary outcome was the rate of survival to 30 days post-arrest. A secondary outcome included the rate of survival until hospital discharge with favourable neurologic outcome (on the modified Rankin scale)

Results

At 30 days 3.2% in the adrenaline group and 2.4% in the placebo group were alive. There was no significant difference in the proportion of patients who survived with favourable outcomes in each group. At the time of hospital discharge, however, severe neurologic impairment had occurred in 31% of the patients in the adrenaline group as against only 17.8% in the placebo group.

Conclusion

In adults with out-of-hospital cardiac arrest whilst the use of adrenaline resulted in a slightly higher rate of 30-day survival those receiving such treatment resulted in nearly twice the rate of survivors with severe neurologic impairment.

John Long is a retired Senior Police Officer and Past Commonwealth Secretary of the Royal Life Saving Society worldwide and is an international director of The Lifesaving Foundation. He is an RLSS Commonwealth Vice President, RLSS UK Life Governor, International Life Saving Knight, and is Patron of the Rashtriya Life Saving Society of India. He has been involved in lifesaving for over 50 years. He is the 2015 recipient of the International Ireland Medal. John is a Member of the Paramedic 2 Adrenalin Trial Management Committee.