

Spring 4-25-2021

## Too Little pH: How Freshwater Acidification Impacts the Abundance of Macrophytes Consumed by Rusty Crayfish

Lauren Tucker  
Intucke@bgsu.edu

Paul Moore Dr.  
*Bowling Green State University*, pmoore@bgsu.edu

Jay Jones Mr.  
*Bowling Green State University*, hwjones@bgsu.edu

Follow this and additional works at: <https://scholarworks.bgsu.edu/honorsprojects>



Part of the [Integrative Biology Commons](#), and the [Terrestrial and Aquatic Ecology Commons](#)

[How does access to this work benefit you? Let us know!](#)

---

### Repository Citation

Tucker, Lauren; Moore, Paul Dr.; and Jones, Jay Mr., "Too Little pH: How Freshwater Acidification Impacts the Abundance of Macrophytes Consumed by Rusty Crayfish" (2021). *Honors Projects*. 594.  
<https://scholarworks.bgsu.edu/honorsprojects/594>

This work is brought to you for free and open access by the Honors College at ScholarWorks@BGSU. It has been accepted for inclusion in Honors Projects by an authorized administrator of ScholarWorks@BGSU.

Too Little pH: How Freshwater Acidification Impacts the Abundance of Macrophytes Consumed  
by Rusty Crayfish

Lauren Tucker

Honors Project

Submitted to the Honors College  
at Bowling Green State University in partial fulfillment  
of the requirements for graduation with

University Honors May 2021

Paul Moore, PhD, Advisor  
Biology Department

Jay Jones, Advisor  
English Department

## Abstract

Anthropogenic activities such as the burning of fossil fuels result in increased atmospheric carbon dioxide (CO<sub>2</sub>) concentration. High levels of atmospheric CO<sub>2</sub> cause chemical shifts in the carbon cycle. Changes in the carbon cycle due to increased CO<sub>2</sub> levels lead to ocean and freshwater acidification. Freshwater acidification is problematic for species that synthesize their own shells as well as species that use olfaction for decision-making. Rusty crayfish (*Faxonius rusticus*) were subject to simulated freshwater acidification and fed two types of macrophyte, Chara (*Chara braunii*) and Myriophyllum (*Myriophyllum sibiricum*). A series of programming language of R (R Core Team, 2019) indicated that simulated freshwater acidification alters the abundance of macrophytes consumed by rusty crayfish. This study demonstrates that freshwater acidification due to atmospheric levels of CO<sub>2</sub> leads to foraging changes in crayfish.

## Introduction

The atmospheric levels of CO<sub>2</sub> have been steadily increasing as the result of human fossil fuel usage coupled with deforestation. As of 2018, the global average amount of CO<sub>2</sub> was at a record high of 407.4 parts per million. Today, CO<sub>2</sub> levels are higher than any point in the past 800,000 years (IGCCP). Levels of CO<sub>2</sub> have been increasing steadily since the Industrial Revolution due to the increase in prevalence of human activities such as the burning of fossil fuels and deforestation (Bogan, 2009). Records show concentrations of atmospheric CO<sub>2</sub> have increased rapidly since the Industrial Revolution; scientific consensus today says CO<sub>2</sub> emissions are anthropogenic (Cerling, 2005). With the increase of CO<sub>2</sub> levels being continual, there are many extenuating impacts that need to be considered. Excess CO<sub>2</sub> in the atmosphere moves beyond the atmosphere and impacts both the biosphere and hydrosphere (Menon, 2007).

When fossil fuels are combusted, CO<sub>2</sub> is released into the atmosphere, which in turn causes the oceans, and other bodies of water, to uptake CO<sub>2</sub>. This process can cause a phenomenon known as ocean acidification (Gattuso, 2011). The ocean plays a fundamental role in the carbon-cycle (Raven, 2005). Carbon is important, as carbon is a primary component of all forms of life on Earth. Carbon atoms are continually exchanged and recycled as CO<sub>2</sub> between the atmosphere and the Earth, forming the carbon-cycle (Menon, 2007). Regarding the ocean or bodies of freshwater, the exchange of carbon as CO<sub>2</sub> occurs between the Earth's atmosphere and the surface of the body of water (Bolin, 1970). When the carbon molecules and the compounds present in the water interact with one another, chemical reactions take place.

Any CO<sub>2</sub> in the atmosphere is exchanged with the upper layer of any water surface through the process of adsorption (Tarbuck, 2006). As CO<sub>2</sub> enters water ionic bonds form with

water producing  $\text{HCO}_3^-$  (Stirling, 2010). Since  $\text{HCO}_3^-$  is a weak acid, hydrogen associates and dissociates freely which forms the carbonate system (Zeebe, 2001). The carbonate system is a set of compounds, which include  $\text{H}_2\text{CO}_3$ ,  $\text{HCO}_3^-$ , and  $\text{CO}_3^{2-}$ , produced by the carbonate equilibrium (Lower, 1999). Because the acidic components  $\text{HCO}_3^-$  are weak acids, hydrogen associates and dissociates from the compounds depending upon the pH of the environment and the concentration of both calcium (Ca) and  $\text{CO}_2$  (Lundegard, 1989). As atmospheric concentrations of  $\text{CO}_2$  change, the equilibrium shifts which creates the disturbances in the pH and chemistry of natural bodies of water. Once  $\text{CO}_2$  has solvated into the ocean, carbonic acid ( $\text{H}_2\text{CO}_3$ ) is formed (Adamczyk, 2009). This carbonic acid then dissociates into a hydrogen ion ( $\text{H}^+$ ) and bicarbonate ( $\text{HCO}_3^-$ ) (Likens, 1979). This results in a large concentration of  $\text{H}^+$ , which increases ocean acidity (Burger, 2020). Thus far, the ocean has experienced a 30% increase in acidity, which is equivalent to a 0.1 decrease in pH (Raven, 2005).

A topic involving alterations in  $\text{CO}_2$  and water chemistry is coral bleaching. Within coral reefs, coralline algal communities are fundamental to the production of carbon sediments and binding fragments of carbonate to the structure of the reefs (McLeod, 2013). As pH in these underwater communities decreases, so does the rate of reef calcification due to the lower concentration of  $\text{CO}_3^{2-}$  (Andersson, 2009). While coral bleaching is driven primarily by ocean warming as opposed to ocean acidification, both are caused by anthropogenic  $\text{CO}_2$  (Pörtner, 2008). In the case of ocean warming, higher concentrations of atmospheric  $\text{CO}_2$  result in periods, lasting between weeks and months, of a locational increase in water temperature (Berkelmans, 2004). The combined impacts of ocean acidification and warming can be detrimental for coral reef systems, amongst others.

Freshwater acidification is also an issue, but freshwater acidification is not subject to as many studies or discussions as ocean acidification. Acidification of inland freshwaters via anthropogenic or ecological sources, such as acid rain, has the potential to be a major stressor for these ecosystems (Hasler, 2018). In areas with a low capacity to neutralize acid such as mountainous regions with high elevation, areas with underlying bedrock, and farmland, acid rain is a common issue (Lajewski, 2003). Acid rain occurs when the burning of fossil fuels emits gaseous sulfur dioxide ( $\text{SO}_2$ ) and nitrogen oxides ( $\text{NO}_x$ ) (Driscoll., 2001). Clouds containing particles of  $\text{SO}_2$  and  $\text{NO}_x$  are formed where these particles are transported to via the wind (Parungo, 1987). Acid is then deposited as rain or snow onto the Earth's surface, completing the process that is known as acid rain (Gattuso, 2011). In the United States, New York and Pennsylvania have been subject to acid rain the most (Stoddard, 1999). Western winds carry  $\text{SO}_2$  and  $\text{NO}_x$  from states surrounding the Ohio Valley, where the concentration of these emissions is the highest, then they are converted into sulfuric ( $\text{H}_2\text{SO}_4$ ) and nitric acids ( $\text{HNO}_3$ ) (Eney, 1987). These acids are deposited over New York and Pennsylvania as acid rain (Likens, 1979). Acid rain results in acidification of aquatic ecosystems and poses several problems. One of these problems is the decreasing pH associated with ocean and freshwater acidification, which can be harmful, particularly to calcifying organisms (Orr, 2005).

As the pH within aquatic environments decreases, calcifying organisms, such as crayfish, coral, mussels, and others, face difficulty synthesizing their calcium carbonate ( $\text{CaCO}_3$ ) shells (Orr, 2005). Within ocean environments, the  $\text{CO}_2$  concentration is increasing, while the carbonate ion ( $\text{CO}_3^{2-}$ ) concentration is decreasing (Andersson, 2005). This lower concentration of  $\text{CO}_3^{2-}$  poses a threat to calcifying marine creatures, as without  $\text{CO}_3^{2-}$ , these organisms face difficulty creating biogenic  $\text{CaCO}_3$  (Feeley, 2004). This means these organisms will not be able

to maintain a sustainable external skeleton.  $\text{CaCO}_3$  shells are an important feature of these calcifying organisms, as they are a means of protection against predators (Hofmann, 2010).  $\text{CO}_2$  emissions can result in limited nutrients within freshwater ecosystems, changes in temperature and water connectivity, and desiccation (Beaune, 2018). Changes in temperature are suspected to cause evapotranspiration, which is the transfer of water from the Earth's surface and the atmosphere (Duan, 2019). Evapotranspiration can cause issues with water connectivity as this process results in the excessive drying, or desiccation, and thus a reduction of water present in aquatic ecosystems (McKenney, 1993). Without enough water present, dry areas within streams and other bodies of freshwater can cause the ecosystem to have a disconnect (Sánchez-Carrillo, 2004). Disconnects within the ecosystem could also result in a limitation of nutrients (Sánchez-Carrillo, 2001). These changes within freshwater ecosystems affect the phenological and physiological of organisms living there, as well as migrations and extinctions (Weatherley, 1988). One example of a physiological change that is experienced in organisms subject to freshwater acidification is the loss of olfaction (Munday, 2009).

Given the lack of light in many aquatic environments, organisms tend to rely more heavily on the chemical senses to make ecological decisions (Dodson, 1994). Chemical signals guide aquatic organism's choices on foraging, mating, avoiding predation, and finding shelter (Hay, 2009). Yet, these behavioral actions are threatened by changes in the environment such as lowered pH. As previously stated, when the pH of freshwater ecosystems is lowered, the ecosystem's chemical composition is altered (Lundegard, 1989). An alteration of the chemicals present in an organism's ecosystem can deter their ability to respond to certain chemical signals, which can cause changes in their behavior (Leduc, 2013). Amino acids are present in neutral aqueous solutions as dipolar ions (Kirkwood, 1939). When acid is added the hydrogen ion ( $\text{H}^+$ )

is taken by the carboxyl group (Gu, 1994). The resulting undissociated carboxyl group is formed and when a base is added, a  $H^+$  is released (Tierney, 1988). The  $H^+$  ions released due to the lowering of pH impact the chemoreception of crayfish (Hay, 2010). Low pH levels cause complications within the binding of amino acids to the crayfishes' receptor sites, resulting in crayfish misreading chemical signals and experiencing behavioral changes, such as differences in preference to food stimuli (Allison, 1992).

Crayfish use olfaction to locate and identify environmental food resources, as well as for movement and homing (Kubec, 2019). Crayfish have a chemosensory organ, called a sensilla, located on their antennules (Horner, 2008). Within the olfactory organ, there are olfactory receptor neurons that send signals to the olfactory lobe located in the crayfish's central nervous system (CNS) when the crayfish senses a smell (Blaustein, 1988). Crayfish rely on their olfactory systems throughout their lifespan to find food (Willman, 1994). However, external stimuli can cause the olfactory systems of crayfish to be modified, which can have a negative impact on the ability to locate and identify their food sources (Hay, 2010). Crustaceans such as crayfish are stimulated to perform particular behaviors by sequences of chemical signals (Hay, 2009). When these sequences of chemical signals are disrupted due to an influx of compounds such as  $CO_2$ , crayfish are unable to perform the same behaviors they would otherwise be able to complete (Tierney, 1986). A decrease in pH because of the increase of  $CO_2$  alters the ability of crayfish to olfaction, and thus has the potential to alter their food preferences (Allison, 1992). When considering these impacts, visible differences in food preference of other aquatic creatures is not unforeseeable.

This experiment focuses on how increasingly lower pH levels in freshwater will impact the abundance of macrophytes eaten by rusty crayfish. Aquatic environments where crayfish are



present rely on them for biomass turnover, as they consume the ecosystem's vegetation, animal waste, and corpses (Momot, 1978). As noted, the lowering of pH as a result of CO<sub>2</sub> can impact crayfish beyond shell formation. The goal of this experiment is to determine whether lowering the pH of a crayfish's ecosystem will cause the crayfish to hold a preference toward one macrophyte over another. Two species of macrophyte, Chara (*Chara braunii*) and Myriophyllum (*Myriophyllum sibiricum*), were used in this experiment. Chara is a species of aquatic plant known for their ability to control the clarity and quality of water (Amirnia, 2019). Myriophyllum is a species of aquatic plant known for its roles as a stabilizer of sediment and a potential detoxifier (Hanson, 2001). If crayfish favor one macrophyte, the other species of macrophytes could overgrow and cause a chain reaction within the ecosystem. The results of this experiment will help to determine how the abundance of different macrophytes within an environment will differ as ocean acidification occurs.

## **Materials and Methods**

### **Collection and Housing of Animals and Plants**

Sixteen from two (non-reproductive) rusty crayfish (*Faxonius rusticus*, *Cambaridae*) were captured from Maple Bay of Burt Lake in Cheboygan County, Michigan (45.4873°N, 84.7065°W) during the summer of 2019. Animals were transported to the laboratory at Bowling Green State University and housed in a flow through system where they were isolated physically and visually, but not chemically, from other animals. During the trials, crayfish were held in the experimental or control aquaria, both containing peat gravel, the macrophytes, and water. Only healthy individuals with all appendages intact were used in the experiments.

Macrophytes, Chara (*Chara braunii*) and Myriophyllum (*Myriophyllum sibiricum*), were kept in two 40-liter aquaria. Plant health was augmented by providing a source of additional CO<sub>2</sub> and full spectrum lights. Additional CO<sub>2</sub> was created by a by-product of yeast fermentation in a system with one 590-milliliter bottle and a 390-milliliter bottle. The first bottle, containing the yeast and fermentation source (pure sugar), was connected via small diameter tigon tubing to the second bottle, which served as collecting spot for yeast and liquid that came through the tubing. The second bottle had an additional tube that provided CO<sub>2</sub> to the holding tanks for the macrophytes.

### **Acidification Impact on Crayfish's Food Preference**

#### **Experimental Design**

To test the effects of freshwater acidification on foraging preferences and consumption of macrophytes, sixteen crayfish were placed within a testing aquarium and provided known quantities of macrophytes to consume over a 5-day period. All crayfish and macrophytes were placed in the test aquaria starting at a pH of 7. To simulate changes in acidification, the pH of the test aquarium was lower by 1 pH unit every day to a final level of pH 4 on day 4. The control aquarium remained at a pH of 7 throughout the experimental period.

To begin the feeding trials, plant samples were then selected from their respective aquaria, dried via salad spinner (Farberware Basics, Item No. 5158683), then weighed to the nearest 0.001 g. These macrophytes were then attached to glass stirring rods (255 x 6 mm: 1 x OD) with 26-gauge green painted floral wire. To ensure the plants would be kept in place for the entire trial, the rods were placed into three hardware cloth brackets (24 x 19 cm: 1 x w). The

arrangement of the macrophytes on the brackets were rotated across trials, as to prevent bias based on location within the aquaria.

### Protocol of Experiment

The experiment began October 2, 2020 and was concluded on December 7, 2020 after completing 8 of the 5-day trials. On day 1, the macrophytes in both experimental and control aquaria were dried, weighed and the mass of the macrophytes would be recorded. The pH in both aquaria were measured and altered until the water in the aquaria reached a unit of 7. The pH of the experimental aquarium, on day 2, would be lowered to 6 via pH down. pH down, composed of phosphoric acid, is produced by General Hydroponics as an acidic buffering solution for pH stability. Before the pH in the experimental aquarium was lowered to 6, the macrophytes in both aquaria were dried, weighed, and their masses were recorded. The macrophyte stems were dried, weighed, attached to the rods, and connected to the bracket in the same position as before, and placed into the experimental and control aquaria. The macrophytes were removed from both aquaria, dried, weighed, and had their masses recorded on day 3. The pH was lowered in the experimental aquarium to 5 after new, dried, and weighed macrophyte stems were added. On day 4, the pH was lowered to the final unit of 4 in the experimental aquarium after removing, drying, and weighing the macrophyte stems in the aquaria, and adding new stems to the glass rods and brackets. Day 5 was the last day of the trial, and after the macrophyte stems were dried and weighed, results would be collected and analyzed.

### **Data Analysis**

All data analysis took place within the programming language of R (R Core Team, 2019). The beginning steps of the data conditioning follow those typically done in mixed model analysis (Zuur, 2009). The first step in the analysis was to produce histograms, qqplots, and normality tests of all the behavioral variables. All data included in the analysis were normally distributed.


All statistical models were performed using generalized linear mixed models (Zuur et al., 2009). All models run in R used the lmer function from the lmerTest package in R (Kuznetsova et al. 2017, R Core Team 2019). Following model construction, the outputs were extracted using the anova function from the car package in R (Fox et al., 2012). For the models, the amount of macrophyte consumed as the dependent variable and the treatment (lowered or constant pH), day (1-4), pH, and macrophyte served as categorical independent measures. Finally, the trial number was used as a random factor within each model.

## **Results**


### **Macrophyte Consumption**

The overall impact of differing pH on the macrophyte consumption of crayfish was found to be significant ( $F_{[3,111,0.05]} = 12.220, p < 0.0007$ ). The linear mixed model for the daily macrophyte consumption of crayfish shows that the consumption of Myriophyllum is affected more by altering pH than the consumption of Chara. When comparing the biomass changes within macrophyte species across treatments, the crayfish consumed much less Myriophyllum than Chara. The amount of Myriophyllum consumed decreased from day 1 to day 3, and then increased on day 4 for both the crayfish receiving treatment and the crayfish in the control group. For the Chara, both groups experienced some variance, but not to the same extent as the Myriophyllum.

These graphs display the distribution of macrophytes consumed by crayfish throughout this experiment. The first graph shows the distribution of Chara consumed, and the second graph shows the distribution of Myriophyllum consumed. The control group experienced an increase in consumption of Chara from day 1 to day 4, with the net increase being approximately 0.02g. The group receiving treatment experienced a greater decrease in Chara consumption from day 1 to day 4, as the difference between these days is approximately 0.08g. For Myriophyllum, the control group and the group receiving treatment experienced decrease in consumption from day 1 to day 3, ending with an increase in consumption from day 3 to day 4. However, the group receiving treatment experienced a much greater increase in the consumption of Myriophyllum than the control group between days 3 and 4. The control group displayed a net decrease of approximately 0.05g in the consumption of Myriophyllum. The group receiving treatment experienced net increase of approximately 0.03g in Myriophyllum consumption. The largest difference in the distribution of macrophytes eaten by crayfish is experienced on day 4. Overall, the crayfish consumed a larger amount of Chara than Myriophyllum, regardless of being under control, or receiving treatment.



**Figure 1.** Distribution of Chara Biomass Consumed by Crayfish.



**Figure 2.** Distribution of Myriophyllum Biomass Consumed by Crayfish.

## Discussion

The results of this experiment show that the consumption of Myriophyllum by crayfish is impacted more than the consumption of Chara. When analyzing all treatments, the crayfish in this study consumed much less Myriophyllum than Chara. The control group experienced a net increase of approximately 0.02 g in Chara consumption and a net decrease of approximately 0.05 g in Myriophyllum consumption. The group receiving treatment experienced a net decrease of approximately 0.08 g in Chara consumption and a net increase of approximately 0.03 g in Myriophyllum consumption. Regardless of the conditions, the crayfish consumed a larger amount of Chara than Myriophyllum. The changes in macrophyte consumption by crayfish are due to the simulation of freshwater acidification performed in this experiment.

Freshwater acidification results in changes within natural processes taking place in the environment. One of these disturbances is a loss of olfaction within crayfish (Munday, 2009). Freshwater acidification causes changes in the perception of chemical signals by crayfish (Hay, 2010). Crayfish use olfactory senses to differentiate between various chemical signals within their environment to make decisions on foraging, mating, and homing (Kubec, 2019). Crayfish are stimulated to perform these behaviors when they receive a sequence of chemical cues or signals (Hay, 2009). To differentiate between chemical signals, crayfish use the sensilla located on their antennules as a chemosensory organ (Horner, 2008). When crayfish sense a smell, the olfactory receptor neurons within their sensilla send a signal to the olfactory lobe in the crayfish's central nervous system (Blaustein, 1988). External stimuli such as freshwater acidification cause modifications within the crayfish's olfactory system due to changes in chemical signals, leading to alterations in the crayfishes' behavior (Leduc, 2013). Low pH levels due to an increasing concentration of CO<sub>2</sub> results in changes in the olfactory systems of crayfish,

which can cause their food preference to change (Allison, 1992). When crayfish experience the effects of freshwater acidification their preference in macrophytes can alter biomass turnover within their habitat (Momot, 1978).

Changes in macrophyte consumption by crayfish can cause differences in biomass turnover than previously experienced in their habitat (Momot, 1978). If one macrophyte is favored over another within an ecosystem relative to their consumption by other species, the less favored macrophyte could overgrow. One macrophyte being preferred over another could cause a chain reaction within their habitat, as each macrophyte species plays a specific role (Rejmankova, 2011). Macrophytes serve many purposes within their habitats, such as influencing the movement of water and providing sustenance as well as shelter for other organisms (Dvořák, 1982). These functions of macrophytes within freshwater ecosystems have influence upon the nutrient cycling and food web interactions that occur (Ziegler, 2015). Changes in the concentration of macrophytes within a habitat can impact the availability for nutrients needed by other organisms, thus altering food web interaction (Warfe, 2006). Differences in these interactions could put stress upon the environment in the form of lower nutrient availability and lower rates of productivity (Cronin, 2003). Lower nutrient availability and productivity alter the processes that would normally occur, especially if the species has a large influence within their ecosystem (Small, 2011). Species with disproportional influences on their environment relative to abundance are known as ecosystem engineers or keystone species (Power, 1996).

Crayfish are commonly referred to as ecosystem engineers or keystone species because they have a disproportionately large effect on their environment relative to their abundance (Creed, 2004). Crayfish are considered keystone species or ecosystem engineers due to their



ability to withstand a broad range in temperature, indicate water quality, and dictate foodweb interactions (Reynolds, 2013). As keystone species, changes in crayfish behavior can result in widespread consequences for other species interactions in their ecosystem (Hart, 1992). Crayfish consume a large amount of macrophytes and other ecosystem waste within their habitat (Momot, 1978). In addition to having an impact on the concentration of macrophytes within their habitat, crayfish can also influence the presence of other species due to the availability of particular macrophytes (Nyström 2002, Reynolds and Souty-Grosset 2012). The change in macrophyte consumption shown by crayfish who have experienced freshwater acidification can impose widespread consequences within their ecosystem.

While this experiment studied the impact of freshwater acidification on the macrophyte consumption of crayfish, other species interactions are altered as well. The crayfish used in this study consumed less *Myriophyllum* than *Chara* across all treatments. For *Chara* consumption, the control group experienced a net increase while the group receiving treatment experienced a net decrease. The opposite occurred for the consumption of *Myriophyllum* as the control group experienced a net decrease while the group receiving treatment experienced a net increase. These changes in macrophyte consumption can be attributed to the impact of simulated freshwater acidification on the olfactory systems of crayfish. The chemical signals received by the crayfish's olfactory systems are altered due to freshwater acidification (Tierney, 1988). Without being able to receive the proper chemical signals crayfish experience behavioral changes such as differences in their eating habits (Allison, 1992). Differences in macrophyte consumption by crayfish can cause problems within their habitat as macrophytes play roles within the environment in terms of nutrient cycling and availability, along with food web interaction (Warfe, 2006). If changes in macrophyte consumption by crayfish occur, ecosystems can

experience lower rates of nutrient production and availability, putting stress upon the ecosystem (Cronin, 2003). This issue is particularly troublesome as crayfish are a keystone species, meaning crayfish are an organism which have a large impact on their environment disproportionate to their abundance (Reynolds, 2013). This study found the reduction of pH as a result of simulated freshwater acidification has an impact on the abundance of macrophytes crayfish consume. Crayfish and the macrophytes they consume play a large role in their ecosystems' nutrient availability and species diversity (Reynolds, 2011). Studies on freshwater acidification are important for environmental sustainability.

## Literature Cited

- Adamczyk, K., Prémont-Schwarz, M., Pines, D., Pines, E., and Nibbering, E. "Real-time observation of carbonic acid formation in aqueous solution." *Science* 326.5960 (2009): 1690-1694.
- Allison, V., Dunham, D., and Harvey, H. "Low pH alters response to food in the crayfish *Cambarus bartoni*." *Canadian Journal of Zoology* 70.12 (1992): 2416-2420.
- Amirnia, S., Asaeda, T., Takeuchi, C., and Kaneko, Y. "Manganese-mediated immobilization of arsenic by calcifying macro-algae, *Chara braunii*." *Science of The Total Environment* 646 (2019): 661-669.
- Andersson, A., Mackenzie, F., and Lerman, A. "Coastal ocean and carbonate systems in the high CO<sub>2</sub> world of the Anthropocene." *American Journal of Science* 305.9 (2005): 875-918.
- Andersson, A., Kuffner, I., Mackenzie, F., Jokiel, P., Rogers, K., and Tan, A. "Net loss of CaCO<sub>3</sub> from a subtropical calcifying community due to seawater acidification: mesocosm-scale experimental evidence." *Biogeosciences* 6.8 (2009): 1811-1823.
- Beaune, D., Sellier, Y., Luquet, G., and Grandjean, F. "Freshwater acidification: an example of an endangered crayfish species sensitive to pH." *Hydrobiologia* 813.1 (2018): 41-50.
- Berkelmans, R., De'ath, G., Kininmonth, S., and Skirving, W. "A comparison of the 1998 and 2002 coral bleaching events on the Great Barrier Reef: spatial correlation, patterns, and predictions." *Coral reefs* 23.1 (2004): 74-83.
- Blaustein, D., Derby, C., Simmons, R., and Beall, A. "Structure of the brain and medulla terminalis of the spiny lobster *Panulirus argus* and the crayfish *Procambarus clarkii*, with an emphasis on olfactory centers." *Journal of Crustacean Biology* 8.4 (1988): 493-519.
- Bogan, R., Ohde, S., Arakaki, T., Mori, I., and McLeod, C. "Changes in rainwater pH associated with increasing atmospheric carbon dioxide after the industrial revolution." *Water, air, and soil pollution* 196.1 (2009): 263-271.
- Bolin, B. "The carbon cycle." *Scientific American* 223.3 (1970): 124-135.
- Burger, F., John, J., and Frölicher, T. "Increase in ocean acidity variability and extremes under increasing atmospheric CO<sub>2</sub>." *Biogeosciences* 17.18 (2020): 4633-4662.
- Cerling, T., Dearing, M., and Ehleringer, J. *A history of atmospheric CO<sub>2</sub> and its effects on plants, animals, and ecosystems*. Springer, 2005.
- Comeau, S., Carpenter, R., and Edmunds, P. "Coral reef calcifiers buffer their response to ocean acidification using both bicarbonate and carbonate." *Proceedings of the Royal Society B: Biological Sciences* 280.1753 (2013): 20122374.

- Creed Jr, R. and Reed, J. "Ecosystem engineering by crayfish in a headwater stream community." *Journal of the North American Benthological Society* 23.2 (2004): 224-236.
- Cripps, I., Munday, P., and McCormick, M. "Ocean acidification affects prey detection by a predatory reef fish." *Plos one* 6.7 (2011): e22736.
- Cronin, G., and Lodge, D. "Effects of light and nutrient availability on the growth, allocation, carbon/nitrogen balance, phenolic chemistry, and resistance to herbivory of two freshwater macrophytes." *Oecologia* 137.1 (2003): 32-41.
- Denissenko, P., Lukaschuk, S., and Breithaupt, T. "The flow generated by an active olfactory system of the red swamp crayfish." *Journal of Experimental Biology* 210.23 (2007): 4083-4091.
- Dodson, S., Crowl, T., Peckarsky, B., Kats, L., Covich, A., and Culp, J. "Non-visual communication in freshwater benthos: an overview." *Journal of the North American Benthological Society* 13.2 (1994): 268-282.
- Doney, S., Fabry, V., Feely, R., and Kleypas, J. "Ocean acidification: the other CO<sub>2</sub> problem." *Annual review of marine science* 1 (2009): 169-192.
- Driscoll, C., Lawrence, G., Bulger, A., Butler, T., Cronan, C., Eagar, C., Lambert, K., Likens, G., Stoddard, J., and Weathers, K. "Acidic Deposition in the Northeastern United States: Sources and Inputs, Ecosystem Effects, and Management Strategies: The effects of acidic deposition in the northeastern United States include the acidification of soil and water, which stresses terrestrial and aquatic biota." *BioScience* 51.3 (2001): 180-198.
- Duan, K., Caldwell, P., Sun, G., McNulty, S., Zhang, Y., Shuster, E., Liu, B., and Bolstad, P. "Understanding the role of regional water connectivity in mitigating climate change impacts on surface water supply stress in the United States." *Journal of Hydrology* 570 (2019): 80-95.
- Dvořák, J., and Best, E. "Macro-invertebrate communities associated with the macrophytes of Lake Vechten: structural and functional relationships." *Studies on Lake Vechten and Tjeukemeer, The Netherlands*. Springer, Dordrecht, 1982. 115-126.
- Eney, A., and Petzold, D. "The problem of acid rain: an overview." *Environmentalist* 7.2 (1987): 95-103.
- Feely, R., Sabine, C., Lee, K., Berelson, W., Kleypas, J., Fabry, V., and Millero, F. "Impact of anthropogenic CO<sub>2</sub> on the CaCO<sub>3</sub> system in the oceans." *Science* 305.5682 (2004): 362-366.
- Gattuso, J., and Hansson, L. *Ocean acidification*. Oxford University Press, 2011.
- Gu, Zhengtian, Zambrano, R., and McDermott, A. "Hydrogen bonding of carboxyl groups in solid-state amino acids and peptides: comparison of carbon chemical shielding, infrared frequencies, and structures." *Journal of the American Chemical Society* 116.14 (1994): 6368-6372.

- Hanson, M., Sibley, P., Mabury, S., Muir, D., and Solomon, K. "Chlorodifluoroacetic acid fate and toxicity to the macrophytes *Lemna gibba*, *Myriophyllum spicatum*, and *Myriophyllum sibiricum* in aquatic microcosms." *Environmental Toxicology and Chemistry: An International Journal* 20.12 (2001): 2758-2767.
- Hart, D. "Community organization in streams: the importance of species interactions, physical factors, and chance." *Oecologia* 91.2 (1992): 220-228.
- Hasler, C., Jeffrey, J., Schneider, E., Hannan, K., Tix, J., and Suski, C. "Biological consequences of weak acidification caused by elevated carbon dioxide in freshwater ecosystems." *Hydrobiologia* 806.1 (2018): 1-12.
- Hay, M. "Marine chemical ecology: chemical signals and cues structure marine populations, communities, and ecosystems." *Annual review of marine science* vol. 1 (2009): 193-212. doi:10.1146/annurev.marine.010908.163708.
- Hay, M. "Crustaceans as powerful models in aquatic chemical ecology." *Chemical communication in crustaceans*. Springer, New York, NY, 2010. 41-62.
- Hofmann, G., Barry, J., Edmunds, P., Gates, R., Hutchins, D., Klinger, T., and Sewell, M. "The effect of ocean acidification on calcifying organisms in marine ecosystems: an organism-to-ecosystem perspective." *Annual review of ecology, evolution, and systematics* 41 (2010): 127-147.
- Horner, A., Schmidt, M., Edwards, D., and Derby, C. "Role of the olfactory pathway in agonistic behavior of crayfish, *Procambarus clarkii*." *Invertebrate Neuroscience* 8.1 (2008): 11-18.
- Kirkwood, J. "Theoretical Studies upon Dipolar Ions." *Chemical Reviews* 24.2 (1939): 233-251.
- Kubec, J., Kouba, A., and Buřič, M. "Communication, behaviour, and decision making in crayfish: a review." *Zoologischer Anzeiger* 278 (2019): 28-37.
- Lajewski, C., Mullins, H., Patterson, W., and Callinan, C. "Historic calcite record from the Finger Lakes, New York: impact of acid rain on a buffered terrane." *Geological Society of America Bulletin* 115.3 (2003): 373-384.
- Leduc, A., Munday, P., Brown, G., Ferrari, M. "Effects of acidification on olfactory-mediated behaviour in freshwater and marine ecosystems: a synthesis." *Philosophical Transactions of the Royal Society B: Biological Sciences* 368.1627 (2013): 20120447.
- Likens, G., Wright, R., Galloway, J., and Butler, T. "Acid rain." *Scientific American* 241.4 (1979): 43-51.
- Lower, S. "Carbonate equilibria in natural waters." Simon Fraser University 544 (1999).
- Lundegard, P., and Land, L. "Carbonate equilibria and pH buffering by organic acids—response to changes in pCO<sub>2</sub>." *Chemical Geology* 74.3-4 (1989): 277-287.
- McLeod, E., Anthony, K., Andersson, A., Beeden, R., Golbuu, Y., Kleypas, J., Kroeker, K., Manzello, D., Salm, R., Schuttenbery, H., and Smith, J. "Preparing to manage coral reefs

- for ocean acidification: lessons from coral bleaching." *Frontiers in Ecology and the Environment* 11.1 (2013): 20-27.
- McKenney, M., and Rosenberg, N. "Sensitivity of some potential evapotranspiration estimation methods to climate change." *Agricultural and Forest Meteorology* 64.1-2 (1993): 81-110.
- Menon, S., Denman, K., Brasseur, G., Chidthaisong, A., Ciais, P., Cox, P., Dickinson, R., Hauglustaine, D., Heinze, C., Holland, E., Jacob, D., Lohmann, U., Ramachandran, S., Wofsy, S., and Zhang, X. *Couplings between changes in the climate system and biogeochemistry*. No. LBNL-464E. Lawrence Berkeley National Lab. (LBNL), Berkeley, CA (United States), 2007.
- Momot, W., Gowing, H., and Jones, P. "The dynamics of crayfish and their role in ecosystems." *American Midland Naturalist* (1978): 10-35.
- Munday, P., Dixson, D., Donelson, J., Jones, G., Pratchett, M., Devitsina, G., and Doving, K. "Ocean acidification impairs olfactory discrimination and homing ability of a marine fish." *Proceedings of the National Academy of Sciences* 106.6 (2009): 1848-1852.
- Orr, J., Fabry, V., and Yool, A. "Anthropogenic ocean acidification over the twenty-first century and its impact on calcifying organisms." *Nature* 437.7059 (2005): 681-686.
- Parungo, F., Nagamoto, C. and Maddl, R. "A study of the mechanisms of acid rain formation." *Journal of Atmospheric Sciences* 44.21 (1987): 3162-3174.
- Pörtner, H. "Ecosystem effects of ocean acidification in times of ocean warming: a physiologist's view." *Marine Ecology Progress Series* 373 (2008): 203-217.
- Power, M., Tilman, D., Estes, J., Menge, B., Bond, W., Mills, L., Daily, G., Castilla, J., Lubchenco, J., and Paine, R. "Challenges in the quest for keystones: identifying keystone species is difficult—but essential to understanding how loss of species will affect ecosystems." *BioScience* 46.8 (1996): 609-620.
- Raven, J., Caldiera, K., Elderfield, H., Hoegh-Guldberg, O., Liss, P., Riebesell, U., Shepherd, J., Turvey, C., and Watson, A. "Ocean acidification due to increasing atmospheric carbon dioxide." *The Royal Society*, 2005.
- Rejmankova, E. "The role of macrophytes in wetland ecosystems." *Journal of Ecology and Environment* 34.4 (2011): 333-345.
- Reynolds, J., and Souty-Grosset, C. *Management of freshwater biodiversity: crayfish as bioindicators*. Cambridge University Press, 2011.
- Reynolds, J., Souty-Grosset, C., and Richardson, A. "Ecological roles of crayfish in freshwater and terrestrial habitats." *Freshwater Crayfish* 19.2 (2013): 197-218.
- Sánchez-Carrillo, S., and Álvarez-Cobelas, M. "Nutrient dynamics and eutrophication patterns in a semi-arid wetland: the effects of fluctuating hydrology." *Water, air, and soil pollution* 131.1 (2001): 97-118.

- Sánchez-Carrillo, S., Angeler, D., Sanchez-Andres, R., Alvarez-Cobelas, M., and Garatuza-Payan, J. "Evapotranspiration in semi-arid wetlands: relationships between inundation and the macrophyte-cover: open-water ratio." *Advances in Water Resources* 27.6 (2004): 643-655.
- Small, G., Pringle, C., Pyron, M., and Duff, J. "Role of the fish *Astyanax aeneus* (Characidae) as a keystone nutrient recycler in low-nutrient Neotropical streams." *Ecology* 92.2 (2011): 386-397.
- Stirling, A., and Papai, I. "H<sub>2</sub>CO<sub>3</sub> Forms via HCO<sub>3</sub><sup>-</sup> in Water." *The Journal of Physical Chemistry B* 114.50 (2010): 16854-16859.
- Stoddard, J., Jeffries, D., Lukewille, A., Clair, T., Dillon, P., Driscoll, C., Forsius, M., Johannessen, M., Kahl, J., Kellogg, J., Kemp, A., Mannio, J., Monteith, D., Murdoch, P., Patrick, S., Rebsdorf, A., Skjelkvale, B., Stainton, M., Traaen, T., Van Dam, H., Webster, K., Wieting, J., and Wilander, A. "Regional trends in aquatic recovery from acidification in North America and Europe." *Nature* 401.6753 (1999): 575-578.
- Tarback, T., and Richmond, G. "Adsorption and Reaction of CO<sub>2</sub> and SO<sub>2</sub> at a Water Surface." *Journal of the American Chemical Society* 128.10 (2006): 3256-3267.
- Tierney, A., and Atema, J. "Effects of acidification on the behavioral response of crayfishes (*Orconectes virilis* and *Procambarus acutus*) to chemical stimuli." *Aquatic toxicology* 9.1 (1986): 1-11.
- Tierney, A., and Telle Atema, T. "Amino acid chemoreception: effects of pH on receptors and stimuli." *Journal of chemical ecology* 14.1 (1988): 135-141.
- Warfe, D., and Barmuta, L. "Habitat structural complexity mediates food web dynamics in a freshwater macrophyte community." *Oecologia* 150.1 (2006): 141-154.
- Willman, E., Hill, A., and Lodge, D. "Response of three crayfish congeners (*Orconectes* spp.) to odors of fish carrion and live predatory fish." *American Midland Naturalist* (1994): 44-51.
- Weatherley, N. "Liming to mitigate acidification in freshwater ecosystems: a review of the biological consequences." *Water, Air, and Soil Pollution* 39.3 (1988): 421-437.
- Zeebe, R., and Wolf-Gladrow, D. *CO<sub>2</sub> in seawater: equilibrium, kinetics, isotopes*. No. 65. Gulf Professional Publishing, 2001.
- Ziegler, J., Solomon, C., Finney, B., and Gregory-Eaves, I. "Macrophyte biomass predicts food chain length in shallow lakes." *Ecosphere* 6.1 (2015): 1-16.